



Weed Diversity and their Management in Major Rice-based Cropping Systems in India

Saptashree Das[✉], Manabendra Ray¹, Aditi Saha Roy¹ and Subhajit Barat²

¹Dept. of Agronomy, Faculty of Agriculture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal (741 252), India

²Dept. of Agronomy, Palli Siksha Bhavana (Institute of Agriculture), Visva-Bharati, Sriniketan, Birbhum, West Bengal (731 236), India



Corresponding ✉ saptashree77@gmail.com

ID 0009-0009-9276-5660

ABSTRACT

Weeds are unwanted and undesirable plants that interfere with cultivated crops for utilization of natural resources such as nutrients, water, and light, space, causing harbor of insect pests and disease adverse allelopathic effects and increasing cost of production indirectly, reducing crop yield. India is an agrarian country, where cropping systems form the backbone of Indian agriculture. Every cropping system has its associated weeds, and to manage these weeds requires a specific crop management approach. The main rice-based cropping systems in India, such as rice-wheat, rice-rice, rice-pulse, rice-maize, and rice-oilseed, have been pivotal in ensuring food security. However, these systems face significant challenges due to weed infestations, leading to yield losses ranging from 13% to 97%. Weeds like grass, sedge and broad-leaf weed varieties pose a substantial threat to these crops, particularly in direct-seeded conditions. To manage weeds effectively, various approaches are employed, including crop rotation, tillage, herbicide application, and intercropping. For instance, in rice-wheat systems, zero-tillage practices, crop residue mulch, and early sowing help to reduce weed emergence. In rice-pulse systems, pre-emergence herbicides, hand weeding, and intercropping prove effective. Maize-based systems benefit from tillage, mulching, and planting at closer spacing. Weed management is critical to sustaining rice-based cropping systems and ensuring food security in India. Tailored strategies that consider crop-specific challenges are essential to mitigate yield losses and maintain agricultural productivity.

KEYWORDS: Rice-based cropping system, herbicide, weed management, yield

Citation (VANCOUVER): Das et al., Weed Diversity and their Management in Major Rice-based Cropping Systems in India. *International Journal of Bio-resource and Stress Management*, 2024; 15(7), 01-09. [HTTPS://DOI.ORG/10.23910/1.2024.5354](https://doi.org/10.23910/1.2024.5354).

Copyright: © 2024 Das et al. This is an open access article that permits unrestricted use, distribution and reproduction in any medium after the author(s) and source are credited.

Data Availability Statement: Legal restrictions are imposed on the public sharing of raw data. However, authors have full right to transfer or share the data in raw form upon request subject to either meeting the conditions of the original consents and the original research study. Further, access of data needs to meet whether the user complies with the ethical and legal obligations as data controllers to allow for secondary use of the data outside of the original study.

Conflict of interests: The authors have declared that no conflict of interest exists.

RECEIVED on 07th April 2024

RECEIVED in revised form on 17th June 2024

ACCEPTED in final form on 28th June 2024

PUBLISHED on 05th July 2024

1. INTRODUCTION

Rice (*Oryza sativa* L.) is the primary cereal crop grown in more than 100 countries worldwide, serving as the fundamental dietary staple where about 90% is produced and consumed for over half of the global population (Rao et al., 2007). India is the world's 2nd highest producer of rice with a production of 130.29 mt from a cultivated area of 46.38 mha. Rice-based cropping system plays a pivotal role in food production, featuring rice as the primary crop. The higher productivity and stability of the rice-based cropping system provide ample employment and income generation opportunities to rural masses besides ensuring food security. To satisfy the needs of a growing population, India must focus on enhancing crop productivity. This can be achieved by identifying and addressing the obstacles that impede farmers from achieving high yields. In India, weed is one of the major biological constraints that limit crop productivity, input-use efficiency and profitability in rice-based cropping systems. Continuous cultivation of rice-wheat cropping sequence favoured the intensification of grassy weeds (Bhatt et al., 2016). Monocropping system favours certain noxious weed species to become dominant in a particular cropping system (Chauhan et al., 2012). They compete with crops for natural resources i.e. light, nutrients, water, and space besides being responsible for reducing the quantity and quality of agricultural productivity (Rao and Nagamani, 2013; Rao et al., 2014.). Weeds are undesired and unwanted plants that negatively impact on human welfare by interfering with the harbor of insect-pests and disease causing organisms, agents causing adverse allelopathy effect (Rao, 2014). Weeds are one of the major issues in agricultural production because of the rise in woody species (Ghersa et al., 2002), herbicide-tolerant species (Hyvonen and Salonen, 2002), and herbicide-resistant biotypes (Heap, 2020) and also because of the negative impact on biodiversity (Satorre et al., 2020) and environmental pollution (Hunt et al., 2017). Among rice based cropping system rice-rice, rice-wheat, rice-maize, rice-pulse, rice-jute, rice-wheat-pulse, rice-oilseed, rice-mustard, rice- pulse/oilseed as paira cropping/utera (lathyrus/lentil/mustard), rice-potato-summer rice are predominant. Each cropping system comes with its unique weed flora, biodiversity and management challenges, necessitating tailored crop management strategies to effectively control these unwanted plants with robust Crop management practices like crop rotation, tillage practice, crop residue management, mulching, water management and nutrients management application of agrochemicals like herbicide, pesticide, insecticide and other agricultural practices (Jat et al., 2021; Nath et al., 2022). These management approaches influence weed seed banks, and weed flora (Marshall et al., 2003) regulating weed dynamics process (weed establishment, competition,

dispersion, herbicide resistance) and weed change patterns (Cordeau et al., 2017). The total annual loss caused by weeds is around 37% (Yaduraju et al., 2006). Under farmer's weed management practices, yield losses range between 13.1% to 22.4%, which need to be reduced by improved weed management practices. In global scale, the yield losses from weeds amounts to 40%, which is more than those coming from collective effect of diseases, insect and pests (Hassan et al., 2005). Crucial weeding interventions have to be done during the early stages of the cropping system, soon after sowing, namely in when plants are still weak competitors. By choosing and implementing the right cropping strategies, farmers can significantly reduce weed pressure, minimize the need for synthetic herbicides, and maintain the productivity and sustainability of their fields. Therefore, identification of system specific problematic weeds and controlling by appropriate combination of strategies management through integrated weed management system (cultural, mechanical, chemical and biological methods) to reduce crop specific weed problem to increase agricultural productivity, profitability by indirectly, sustainability, and reduce environmental impacts.

Table 1: Potential yield loss due to weeds in major crops of India (Rao et al., 2014)

Crop	Yield loss (%)	Crop	Yield loss (%)
Chick pea	10-50	Pea	10-50
Cotton	40-60	Pearl millet	16-65
Finger millet	50	Pigeon pea	20-30
Green gram	10-45	Potato	20-30
Groundnut	30-80	Rice	10-100
Horse gram	30	Sorghum	45-69
Jute	30-70	Soybean	10-100
Lentil	30-35	Sugarcane	25-50
Maize	30-40	Vegetables	30-40
Niger	20-30	Wheat	10-60

2. PREDOMINANT WEED FLORA IN RICE-RICE CROPPING SYSTEM

Rice-rice cropping system mainly practices in Uttar Pradesh, Andhra Pradesh, Kerala, Tamilnadu, Chhattisgarh, Punjab, Assam, West Bengal, Orissa, Karnataka, and Gujarat. Due to practice of same crop in same land promote a rich diversity of weed species that may cause yield loss upto 50–90% if weeds are not properly controlled (Chauhan and Johnson, 2011; Singh et al., 2005). It may go up to 97% in upland rice (Singh et al., 1988), 12–70% in transplanted rice and 85% in wet

seeded rice (Singh et al., 2011). Major grassy weeds and sedges commonly associated with rice cultivation include barnyard grass (*Echinochloa colona*), para grass (*Brachiaria mutica*), Chinese sprangletop (*Leptochloa chinensis*), rice flat sedge (*Cyperus iria* L.), purple nutsedge (*Cyperus rotundus*), yellow nut sedge (*Cyperus esculentus*) broad-leaved weeds include horse purselane (*Trianthema portulacastrum* L.), dayflower (*Commelina benghalensis* L.), alligator weed (*Alternanthera philoxeroides*), water primrose (*Ludwigia parviflora*), native gooseberry (*Physalis minima*), pink node flower (*Caesulia axillaris*), and blistering ammania (*Ammania multiflora*). *Cynodon dactylon* poses a significant challenge in fields practicing permanent zero tillage (ZT) within the eastern IGP (Jat et al., 2010). Generally, weed-related issues are more pronounced in direct-seeded rice (DSR) when compared to transplanted conditions (Chhokar et al., 2014). Wild rice (*Echinochloa colona*) is emerging as a major problem in direct-seeded rice (Rao and Nagamani, 2007; Rao et al., 2007). The weeds and the crop emerge at the same time with weeds growing more quickly (Rao et al., 2007), leading up to 50–90% yield reduction if weeds are not timely managed (Chauhan and Johnson, 2011; Singh et al., 2005).

Continuous submergence of the crop successfully controls the weed population and inhibits their germination in transplanted rice fields (Subramanayam et al., 2007). Reports highlighted that the combination of intensive puddling with submergence resulted in the lowest weed dry weight, at 6.63 gm⁻². Practices like soil solarization, planting competitive crop varieties, preparing stale seedbeds, applying mulch, proper fertilization, and practicing intercropping have been found effective for weed management in aerobic rice systems. The rice-sesbania co-culture and killing with 2, 4-D ester at 0.5 kg ha⁻¹ suppress the growth of weeds and is highly effective in controlling both broad-leaved weeds and sedges, but less effective for grassy weeds. However, hand weeding at 20 and 40 DAT gave the highest weed control efficiency with the highest rice grain yield (5.08 t ha⁻¹) (Rekha et al., 2002; Pal et al., 2009). According to Thiagarajan et al. (2002), the use of cono-weeder led to a 10% boost in grain yield during the wet season, whereas it was only 3% higher in the dry season compared to the conventional weeding method. In rice nurseries, the use of effective herbicides such as pretilachlor+safener (Rao and Moody, 1988), cyhalofop-butyl (Jayadeva et al., 2002; Sharma et al., 2004) and propanil in combination with hand-weeding results in healthy rice seedlings for transplanting. Rice ratooning or re-sprouting poses a significant challenge in no-till rice-rice as well as rice-pulses farming systems in eastern Gangetic plains and peninsular India. Application of pyrazosulfuron ethyl @ 20 or 25 g ha⁻¹ at 3 or 10 DAT significantly reduced the weed density and weed dry matter in transplanted rice during the *kharif*

season (Chopra and Chopra, 2003; Shekhar et al., 2004). Bispyribac sodium 10% SC @ 25 g a.i. ha⁻¹ at 15–25 DAS is effective in controlling all three types of weed flora (grasses, broadleaved and sedges). Saha and Rao (2009) recorded the highest weed control efficiency (95.2%) with the application of bensulfuron methyl @ 60 g ha⁻¹ at 20 DAS. Subramanian et al. (2006) noted that higher grain yield (5744 kg ha⁻¹) with the application of pretilachlor with safener+dhaincha intercropping+azolla dual cropping on 30 DAS in wet seeded rice.

3. PREDOMINANT WEED FLORA IN RICE-WHEAT CROPPING SYSTEM

In India, the rice-wheat cropping system is the most prevalent sequence covering around 10.5 million ha mainly in the Indo-Gangetic Plains (IGP) (Jat et al., 2013). The advent of the Green Revolution brought about a significant surge in wheat production in this region. However, the dwarf wheat variety used in this revolution is more susceptible to weed infestation, particularly when nutrient and water resources are abundant. Weeds typically pose a formidable challenge to wheat cultivation, potentially causing 60% reduction in grain yield under conventional tillage and 70% in zero tillage conditions (Jain et al., 2007) if not effectively managed at critical stages. The major weeds commonly found in wheat fields include little seed canary grass (*Phalaris minor*), strawberry clover (*Trifolium fragiferum*), common lamb's quarters (*Chenopodium album*), common vetch (*Vicia sativa*), black nightshade (*Solanum nigrum*) and toothed dock (*Rumex dentatus*) in varying levels of abundance. One strategy to mitigate the weed problem in wheat is to optimize water management during rice cultivation, which can prove beneficial in reducing the presence of weeds like *Rumex*, as they tend to thrive predominantly in wet soil conditions.




Figure 1: Relative weed density and dry weight of weeds in rice-wheat cropping system

It is imperative to implement weed control measures 15–45 days after wheat sowing as this is the critical period for weed interference. The adoption of zero tillage practices and the retention of crop residues on the soil surface have proven to be highly effective methods for reducing the germination

of weeds such as *Phalaris*, *Chenopodium*, *Physalis*, *Anagalis*, and others. For zero-tilled wheat, the use of rice residue mulch @ of 6–10 t ha⁻¹, coupled with early sowing, resulted in the reduction of little seed canary grass emergence by more than 80% (Kumar et al., 2013). Effectively managing water during rice cultivation can also play a beneficial role in decreasing the populations of *Rumex* in wheat, as it primarily thrives in moist soil conditions. Under transplanted rice, application of butachlor @ 1.5 kg ha⁻¹ at 3 DAT followed by 2 hand weeding at 30 and 60 DAS produced significantly higher grain yield than in WSR (wet seeding in puddled soil) with cyhalofopbutyl @ 100 g ha⁻¹ and DSR (dry seeded rice in unpuddled soil) with pendimethalin 1 kg ha⁻¹ under rice-wheat cropping system (Singh et al., 2005). In ZT wheat, pre-plant application of non-selective systemic herbicide glyphosate @ 1 kg a.i. ha⁻¹ at 1–7 DBF effectively controls weeds. Effective control of annual grassy weeds in wheat was achieved by application of post-emergence herbicide clodinafop (Topik) 60 g a.i. ha⁻¹ or sulfosulfuron (Leader) 25 g a.i. ha⁻¹ following the first irrigation at 30 DAS (Gopal et al., 2019). Application of isoproturon+2, 4 D (0.75+0.6 a.i. kg ha⁻¹) produced a higher yield (3.89 t ha⁻¹) and B:C ratio (2.59) in ZT wheat (Singh et al., 2010). A report stated that clodinafop @ 60 g, sulfosulfuron @ 25 g and fenoxaprop-ethyl @ 100 g ha⁻¹ increased the yield of wheat by 41.9, 43.7 and 39.9% respectively over the weedy check (Walia et al., 2006).

4. PREDOMINANT WEED FLORA IN PREDOMINANT WEED FLORA IN RICE-MAIZE CROPPING SYSTEM

Maize-based cropping systems are prevalent on a large scale, particularly in the central and northern parts of the country. The low productivity of maize in India can be attributed to several limiting factors, of which poor weed management poses a major threat to crop productivity. The most important weeds that can be associated with maize-based cropping systems are *Echinochloa colonum*, *Digitaria sanguinalis*, *Dactyloctenium aegyptium*, *Eleusine indica*, *Setaria glauca*, *Sorghum halepense*, *Panicum spp.*, *Cynodon dactylon*, *Digitaria setigera*, among grasses; *Ageratum conyzoides*, *Galinsoga parviflora*, *Commelina benghalensis*, *Undernia cilata*, *Polygonum hydropiper*, *Euphorbia geniculata*, *Oxalis latifolia*, *Celosia argentea*, *Cleome viscosa*, *Sida acuta*, *Aschynomene indica*, *Acanthospermum hispidum*, *Portulaca oleracea*, *Phyllanthus niruri*, *Amaranthus viridis*, *Acalypha indica*, *Tridax procumbens*, *Ipomoea pestigridis*, *Parthenium hysterophorus* and *Euphorbia hirta* among non-grassy weeds and *Cyperus rotundus* and *Cyperus iria* among sedges that reduce the maize yields by 27–60%, depending upon the growth and persistence of weed population (Rout and Staphathy, 1996; Lamani et al., 2000). In the rainy season, it was reported that the emergence of maize and weeds was

simultaneous and the first 20–60 days was the most critical period of competition for the crop (Porwal, 2000; Nayital et al., 1989).

The agronomic practices, viz. tillage and inter-cultivation, intercropping, mulching, cover crops, crop rotation, higher seed rate, planting at closer spacing, nutrient management, planting methods, and other agro-techniques are used for weed management in maize-based cropping systems. Sharma et al. (2000) opined that hoeing at 15 DAS effectively controlled the weed population at 30 DAS which was less than half (23–32 weeds m⁻²) as compared with no intercultural operation (67–70 weeds m⁻²). The intercultural operations like mechanical weeding or two-hand weeding at 20–30 and 35–45 DAS effectively minimized the weed population and increased maize yield (Saikia and Pandey, 2001). Saini et al. (2013) revealed that soybean intercropping +one mechanical weeding (20 DAS) recorded significantly lowest weed dry weight, higher yield attributes and maize equivalent yield which was at par with 2 mechanical weedings (20 and 40 DAS)+mash intercropping in maize. Rani et al. (2011) reported that application of sulfosulfuron 15 g ha⁻¹+imazethapyr 25 g ha⁻¹ as pre-emergence with hand weeding at 40 DAS was found to be effective and economic weed management practice for irrigated sweet corn during Rabi season. Walia et al. (2006) reported that the application of atrazine+alachlor @0.50+0.75 kg ha⁻¹ fb one HW proved to be significantly superior with the highest weed control efficiency of 89.4%.

5. PREDOMINANT WEED FLORA IN RICE-OILSEED CROPPING SYSTEM

The critical period of crop weed competition in groundnut is 40–60 DAS, for sunflower 15–25 DAS, soybean 20–45 DAS and for rapeseed mustard 15–40 DAS. The major weed flora dominated in rice-oilseed cropping system are *Echinochloa colona*, *Digitaria sanguinalis*, *Cynodon dactylon*, *Cyperus rotundus*, *Cyanotis axillaris*, *Ageratum conyzoides*, *Spilanthus paniculata*, *Polygonum plebeium*, *Gnapahalium purpureum*, *Chenopodium album*, *Physalis minima*, *Eclipta alba*, *Cleome viscosa*, *Ludwigia parviflora*, *Solanum nigrum* and *Indigofera hirsute*, *Melilotus indica*, *Trigonella polycerata*, *Chenopodium album*, *Orobancha aegyptiaca*, *Convolvulus arvensis*, *Dactyloctenium aegyptium*, *Digera arvensis*, *Tridax procumbens*, *Phyllanthus niruri*, *Commelina benghalensis*, *Eclipta alba*, *Chenopodium album*, *Parthenium spp* (Mishra et al., 2020; Aruna et al., 2018), yield loss ranges from 13% to 100% depending on the season, cultivars, weed composition and duration of crop weed competition, and the packages of practices adopted (Yaduraju et al., 1980; Kalaiselvan et al., 1994; Rajendran and Louduraj, 1999; and Ghosh et al., 1999). However, Giri et al. (1998) reported that an average yield loss due to weed infestation in irrigated

summer groundnut is 89%. Bunch-type cultivars have minimal leaf cover, horizontal growth habits, late seedling emergence and slow initial growth, and poor competitive abilities; the crop-weed competition is at its highest in the early phases. (Sheoran et al., 2015).

According to Jha and Soni (2013), the flatbed method of sowing recorded a higher density of monocot (43.2 no. m^{-2}) and dicot weeds (34.4 no. m^{-2}) whereas, the broad bed furrow method recorded the lowest weed density (25.1 and 18.7 no. m^{-2}) in soybean. Marangoni et al. (2013) revealed that compared to planting in December, November planted soybean cultivars recorded the lowest dry weight of monocot weeds and total weed dry weight at 26 days after emergence and harvest. The optimum time of sowing enhanced crop development and the competitive potential of soybean against the weed population. According to Singh (2006) practicing several cultural methods like solarization, using transparent or black polyethylene film for 30 days before sowing of soybean, decreased the weed germination. Besides, soil solarization using transparent polyethylene for 30 days increased the soybean yield by 90%. Arora and Tomar (2012) reported the lowest dry weight of weeds, the highest WCE (54%) and the highest pod yield from stale seedbed in groundnut. As per the report of All India Coordinated Research Project on Groundnut (2009) weed population in ground nut was effectively managed by pre-emergence application of pendimethalin @ 1.0 kg ha^{-1} +post-emergence application of quizalofop-p-ethyl @ 50 g ha^{-1} at 20 DAS+one hand weeding at 45 DAS. Application of pre-emergence herbicide pendimethalin @ 1.0 kg ha^{-1} +post-emergence application of imazethapyr @ 75 g ha^{-1} at 20 DAS+one-hand weeding at 45 DAS was proved to be the best weed control in groundnut. Degra et al. (2006) showed that reduction in weed density by pre-planting application of fluchloralin @ 1.0 kg ha^{-1} followed by one-hand weeding at 40 days after sowing in rice under rice-mustard cropping system. According to Singh et al. (2009) pre-emergence application of isoproturon @ 0.5 kg ha^{-1} +one inter-cultivation at 30 DAS brought a significant reduction in weed density and biomass in rapeseed and mustard in India. According to Reddy et al. (1998) application of anilofos @ 0.4 kg ha^{-1} or butachlor @ 1.5 kg ha^{-1} or 2, 4-DEE @ 1.0 kg ha^{-1} at 5 DAT was found equally effective in suppressing weed growth and increasing grain yield of *kharif* rice. During *rabi*, application of oxyfluorfen @ 0.15 kg ha^{-1} and metolachlor @ 1.5 kg ha^{-1} to sunflower was effective in controlling weeds and resulted in a similar yield to that of hand-weeded (20 DAS) crop in the rice-sunflower cropping system.

6. PREDOMINANT WEED FLORA IN RICE-PULSE CROPPING SYSTEM

India is a major pulse producing country in the world sharing about 35% area and 28% production but the area under pulse production has remained static during the past

30 years. Pulses mainly grown as a component crop in the rice-wheat cropping systems in Indo Gangetic Plain (IGP) of India. Continuous practice of rice-wheat cropping system resulted in declining productivity, depletion of soil nutrients and water resources, and frequent outbreaks of pests and diseases, thus emphasizing the need for crop rotation. There is a need to revitalize pulse crops under rice-based cropping system since pulses have the capacity of biological nitrogen fixation (BNF), to improve soil fertility and maintain the long-term sustainability of this production system (Ali and Kumar, 2002). IGP region shows huge potential for expansion of pulses either as a catch crop, sole crop as a component for crop rotation in different cropping systems. Chickpeas (*Cicer arietinum* L.), lentil (*Lens culinaris* L.) and pigeon peas (*Cajanus cajan* L.) are being cultivated as major pulses in IGP of India. Weeds were the major biotic constraints for successful reintroduction and profitable production of pulses (Pande et al., 2000). The initial growth stages of pigeon pea are the most critical stage for weed competition resulting in yield loss up to 90%. Late sowing of chickpeas in a rice-chickpea sequential system exposes chickpeas to greater weed competition. *Chenopodium album* is a major weed of chickpea crop and yield loss can be up to 40%. Lentil is a poor competitor with weed flora due to its slow growth and inadequate weed control may cause yield losses of 20–30%. The predominant weed flora in Rice-pulse cropping system are *Echinochloa colona*, *Cyperus* sp, *Panicum* sp and *Trianthema portulacastrum* (Reddy et al., 2000), *Paspalum paspalodes*, *Marsilea quadrifolia*, *Eclipta alba*, *Ephorbia prostrata* and *Alternanthera triandra*, *Leersia hexandra* and *Echinochloa crusgalli*, *Fimbristylis littoralis* (Pramanick et al., 2014).

In the rice-green gram cropping system application of pre-emergence (PE) herbicide thiobencarb+2, 4-D in rice followed by one-hand weeding (HW) in succeeding crop green gram resulted in the most effective weed control method (Srinivasan et al., 1991). PE application of pendimethalin @ 1 kg ha^{-1} +HW at 30 DAS resulted in lower weed density and higher WCE at all stages of green gram (Hasanain et al., 2020; Rathika et al., 2023). In zero till sown rice fallow black gram cropping system best combination of weed control was dibbling the seeds three days after application of paraquat @ 0.5 kg ha^{-1} with post-emergence application of either fenoxaprop-p-ethyl @ 75 g ha^{-1} or cyhalofop butyl @ 100 g ha^{-1} on 15 DAS resulted in higher grain yield (Sasikala et al., 2014). In the rice-black gram cropping system, the application of anilofos @ 0.4 kg ha^{-1} at 5 DAT effectively suppressed weed growth and increased grain yield in *kharif* rice as that of hand weeding twice at 20 and 40 DAT and the application of pendimethalin, metolachlor, alachlor each @ 1.5 kg ha^{-1} to black gram was equally effective in controlling weeds (Reddy et al., 2000).

Rao et al. (2011) found that post-emergence application of quizalofop ethyl @ 50 g ha⁻¹ recorded higher seed yield, net monetary returns and BCR of rice fallow black gram cropping system. In lentil, PE application of pendimethalin @ 0.75 kg ha⁻¹ fb one hand weeding or imazethapyr @ 25–40 g ha⁻¹ either at 25 or 35 DAS improved the grain yield (1662 kg ha⁻¹) (Yadav et al., 2013; Singh et al., 2014). In the rice-lathyrus cropping system early PoE application of imazosulfuron 10% SC @ 100 g a.i. ha⁻¹ at 5 DAT in rice is the best weed control method resulting in higher grain yield and B:C in both crops (Pramanick et al., 2014). However, application of almix 20 WP @ 4 g ha⁻¹ at 15 DAT also gave good results in the rice-lathyrus cropping system (Dey et al., 2021).

Intercropping enhances crop canopy and thus suppresses weeds. Short-duration legumes, viz. urdbean, mungbean, soybean and cowpea when grown with pigeonpea under an intercropping system suppressed weed flora considerably. The Sorghum+Pigeonpea intercropping system is the most effective weed control cropping system in central and peninsular India. Kaur et al. (2015) reported that the application of PE pendimethalin 1.0 kg ha⁻¹ fb PoE application of imazethapyr 0.10 kg ha⁻¹ at 30–35 DAS has been found effective weed control method in pigeon pea. Talnikar et al. (2008) reported that PE application of alachlor 2 kg ha⁻¹ with HW and hoeing at six weeks after sowing proved most effective and economical in controlling weeds and enhancing the grain yield in pigeon pea + soybean intercropping system.

7. CONCLUSION

Weeds are the most important limiting factors responsible for low yield and hampers crop growth and development. Different rice-based cropping systems have different associated weeds, reduces crop yield and increase disease pest infestation. Therefore, cropping system specific weed management is crucial. An effective cropping system with right crop management, planting, mulching, water management, time of fertilizer application, tillage management etc. not only addresses weed management but also contributes to the overall health and productivity of the agro ecosystem.

8. FUTURE RESEARCH THRUST

Though huge number of research work has already been done in different rice-based cropping systems in different agro climatic zones of the country. The most of the work is confined in the area of different herbicide and mechanical weed control under different crops and cropping systems. Future research work should also focus on special breeding programme for sustainable cultivars of rice and component crops which can suppress the major weeds in

different agro climatic zones. Herbicide resistance is one of the important issues which is to be addressed in future. Many new herbicides are coming very fast and their efficacy should be tested under different resource base.

9. REFERENCES

- Ali, M., Kumar, S., 2002. Diversification in cropping systems through pulses. In: Ali, Chaturvedi, and Gurha, S.N. (Eds.), Pulses for sustainable agriculture and national security. proceedings of national symposium. Indian Institute of Pulses Research, Kanpur, India, Pages 13–22.
- Arora, A., Tomar, S.S., 2012. Effect of soil solarization on weed seed bank in soil. Indian Journal of Weed Science 44(2), 122–123.
- Aruna, E., Sagar, G.K., 2018. Weed management in groundnut under rice-fallow. Indian Journal of Weed Science 50(3), 298–301.
- Bhatt, R., Kukal, S.S., Busari, M.A., Arora, S., Yadav, M., 2016. Sustainability issues on rice-wheat cropping system. International Soil and Water Conservation Research 4, 64–74.
- Chauhan, B.S., Singh, R.G., Mahajan, G., 2012. Ecology and management of weeds under conservation agriculture: A review. Crop Protection 38, 57–65.
- Chauhan, B.S., Johnson, D.E., 2011. Growth response of direct-seeded rice to oxadiazon and bispyribac-sodium in aerobic and saturated soils. Weed Science 59, 119–122.
- Chhokar, R.S., Sharma, R.K., Gathala, M.K., Pundir, A.K., 2014. Effects of crop establishment techniques on weeds and rice yield. Crop Protection 64, 7–12.
- Chopra, N.K., Chopra, N., 2003. Effect of doses and stages of application of pyrazosulfuron ethyl on weeds in transplanted rice. Indian Journal of Weed Science 35(1 and 2), 27–29.
- Cordeau, S., Smith, R.G., Gallandt, E.R., Brown, B., Salon, P., Di Tommaso, A., Ryan, M.R., 2017. Timing of tillage as a driver of weed communities. Weed Science 65, 504–514.
- Dey, R., Sana, M., Pal, S., Banerjee, H., 2021. Efficacy of herbicides against weeds in rice (*Oryza sativa* L.)-lathyrus cropping system. Journal of Crop and Weed 17(1), 293–299.
- Ghersa, C.M., de la Fuente, E.B., Suarez, S.A., Leon, R.J.C., 2002. Woody species invasion to the Rolling Pampa grasslands. Agriculture, Ecosystems & Environment 88(3), 271–278.
- Ghosh, P.K., Mandal, K.G., Kuntal, M.H., 2000. Allelopathic effects of weeds on groundnut (*Arachis hypogaea*) in India-A review. Agricultural Reviews 21, 66–69.

- Giri, A.N., Bhosle, R.H., Shelke, D.K., 1998. Weed management in cotton-groundnut sequence. *Indian Journal of Agronomy* 43, 50–56.
- Gopal, R., Jat, R.K., Malik, R.K., Kumar, V., Alam, M.M., Jat, M.L., Gupta, R., 2019. Direct dry seeded rice production technology and weed management in rice-based systems. *Gates Open Research* 3(207), 207.
- Hassan, G., Khan, I., Khan, H., Munir, M., 2005. Effect of different herbicides on weed density and some agronomic traits of wheat. *Pakistan Journal of Weed Science Research* 11, 17–22.
- Heap, I., 2020. International survey of herbicide resistant weeds. <http://www.weedscience.org>.
- Hunt, N., Hill, J., Liebman, M., 2017. Reducing freshwater toxicity while maintaining weed control, profits, and productivity: effects of increased crop rotation diversity and reduced herbicide usage. *Environmental Science & Technology* 51, 1707–1717.
- Hyvonen, T., Salonen, J., 2002. Weed species diversity and community composition in € cropping practices at two intensity levels-a six-year experiment. *Plant Ecology* 159, 73–81.
- Jain, N., Mishra, J.S., Kewat, M.L., Jain, V., 2007. Effect of tillage and herbicides on grain yield and nutrient uptake by wheat (*Triticum aestivum*) and weeds. *Indian Journal of Agronomy* 52(2), 131–134.
- Jat, H.S., Kumar, V., Kakraliya, S.K., Abdallah, A.M., Datta, A., Choudhary, M., Sharma, P.C., 2021. Climate-smart agriculture practices influence weed density and diversity in cereal-based agri-food systems of western Indo-Gangetic plains. *Scientific reports* 11(1), 15901.
- Jat, M.L., Gathala, M.K., Saharawat, Y.S., Tetarwale, J.P., Gupta, R., Singh, Y., 2013. Double no-till and permanent raised beds in maize-wheat rotation of north-western Indo-Gangetic plains of India: Effects on crop yields, water productivity, profitability and soil physical properties. *Field Crops Research* 149, 291–299.
- Jat, R.K., Gopal, R., Jat, M.L., Saharawat, Y.S., Gupta, R.K., 2010. Conservation agriculture: Strategies for long-term weed management in the eastern Indo-gangetic plains of India. *Conservation Agriculture Course* 63.
- Jayadeva, H.M., Basavarajappa, D.N., Bhairappanavar, S.T., Dinesh, G.R., Shivanna, S., 2002. Evaluation of new herbicide molecules for weed control in wet rice nursery. *Current Research-University of Agricultural Sciences (Bangalore)*, 31, 179–181.
- Jha, A.K., Soni, M., 2013. Weed management by sowing methods and herbicides in soybean. *Indian Journal of Weed Science* 45(4), 250–252.
- Kalaiselvan, P., Manoharan, V., Balasubramanian, T.N., 1994. Effect of different doses of herbicides on the control of groundnut weeds. *Madras Agriculture Journal* 81, 628–630.
- Kaur, R., Raj, R., Das, T.K., Shekhawat, K., Singh, R., Choudhary, A.K., 2015. Weed management in pigeonpea-based cropping systems. *Indian Journal of Weed Science* 47, 267–276.
- Kumar, V., Singh, S., Chhokar, R.S., Malik, R.K., Brainard, D.C., Ladha, J.K., 2013. Weed management strategies to reduce herbicide use in zero-till rice-wheat cropping systems of the Indo-gangetic plains. *Weed Technology* 27(1), 241–254.
- Lamani, C.S., Palled, V.B., Shenoy, H., 2000. Weed dynamics as influenced by integrated nutrient management in hybrid maize. *Current Research* 29(1–2), 21–22.
- Marangoni, R.E., Jakelaitis, A., Tavares, C.J., Rezende, B.P.M., Mello Filho, O.L., Cunha, P.C.R., 2013. Effect of weed interference on soybean cultivars at two sowing times. *Planta Daninha* 31, 511–519.
- Marshall, E.J.P., Brown, V.K., Boatman, N.D., Lutman, P.J.W., Squire, G.R., Ward, L.K., 2003. The role of weeds in supporting biological diversity within crop fields. *Weed Research* 43, 77–89.
- Mishra, J.S., Kumar, R., Mondal, S., Poonia, S.P., Rao, K.K., Dubey, R., Bhaskar, S., 2022. Tillage and crop establishment effects on weeds and productivity of a rice-wheat-mungbean rotation. *Field Crops Research* 284, 108577.
- Mohammad, H., Shukla, D.K., Singh, R.K., Upadhyay, P.K., Shekhawat, K., Rathore, S.S., 2020. Effect of nutrient and weed management on growth and yield of summer mung bean (*Vigna radiata*). *Indian Journal of Agricultural Sciences* 90(4), 771–774.
- Nath, C.P., Hazra, K.K., Kumar, N., Singh, S.S., Praharaj, C.S., Singh, U., Singh, N.P., Nandan, R., 2022. Impact of crop rotation with chemical and organic fertilization on weed seed density, species diversity, and community structure after 13 years. *Crop Protection* 153, 105860.
- Nayital, S.C., Sharma, J., Singh, C.M., 1989. Weed-crop competition studies in maize under Kulu valley conditions. *Indian Journal of Weed Sciences* 21(1–2), 47–50.
- Pal, D., Ghosh, R.K., Sounda, G., Dolai, A.K., Pramanick, M., Mallick, S., 2009. Bio efficacy of some promising herbicides in transplanted *kharif* rice and their influence on soil microflora. *Indian Agriculture* 53(3 and 4), 193–198.
- Pande, S., Sharma, S.B., Rarnakrishna, A., 2000. Biotic stresses affecting legumes production in the Indo-Gangetic Plain. 129–155 In: Johansen, C., Duxbury, J.M., Virmani, S.M., Gowda, C.L.L., Pande, S., Joshi, P.K. (Eds.), *Legumes in rice and wheat cropping*

- systems of the indo-gangetic plain-constraints and opportunities, International Crops Research Institute for the Semi-Arid Tropics, Patancheru, Andhra Pradesh, India and Ithaca Cornell University, New York, USA.
- Porwal, M.K., 2000. Economics of weed-control measures in winter maize (*Zea mays*). Indian Journal of Agronomy 45(2), 433–447.
- Pramanick, B., Bera, P.S., Kundu, C.K., Bandopadhyay, P., Brahmachari, K., 2014. Effect of different herbicides used in transplanted rice on weed management in rice-lathyrus cropping system. Journal of Crop and Weed 10(2), 433–436.
- Rajendran, K., Lourduraj, A.C., 1999. Weed management in groundnut a review. Agricultural Review 20, 59–62.
- Rani, B.S., Sagar, G.K., Reddy, P.M., 2011. Effect of integrated weed management with low volume herbicides in sweet corn (*Zea mays*). Indian Journal of Weed Science 43(1 and 2), 110–112.
- Rao, A.N., Wani, S.P., Ladha, J.K., 2014. Weed management research in India-an analysis of the past and outlook for future 1–26. In: Souvenir (1989–2014). DWR Publication No. 18. Directorate of Weed Research, Jabalpur, India.
- Rao, A.N., Nagamani. A., 2013. Eco-efficient weed management approaches for rice in tropical Asia. Proc. 4th Tropical weed science conf. weed management and utilization in the tropics held during January 23–25, 2013, The Empress Hotel, Chiang Mai, Thailand, 78–87.
- Rao, A.S., 2011. Bio-efficacy of quizalofop ethyl on echinocloa control in rice-fallow black gram. The Andhra Agricultural Journal 58, 130–132.
- Rao, A.N., Nagamani, A., 2007. Available technologies and future research challenges for managing weeds in dry-seeded rice in India. In: Proc. of the 2151 Asian-Pacific Weed Science Society Conf. 2nd–6th October 2007, Colombo, Sri Lanka.
- Rao, A.N., Johnson, D.E., Sivaprasad, B., Ladha, J.K., Mortimer, A.M., 2007. Weed management in direct-seeded rice. Advances in Agronomy 93, 153–255.
- Rao, A.N., Moody, K., 1988. Weed control in rice seedling nurseries. Crop Protection 7, 202–206.
- Rathika, S., Udhaya, A., Ramesh, T., Shanmugapriya, P., 2023. Weed management strategies in green gram: A review. The Pharma Innovation Journal 12(3), 5574–5580.
- Reddy, M.D., Reddy, C.N., Devi, M.P., 2000. Effect of herbicides on weed growth crop performance in rice-blackgram cropping system. Indian Journal of Weed Science 32(3 and 4), 169–172.
- Reddy, C.N., Reddy, M.D., Reddy, N.V., Devi, M.P., 1998. Weed management in rice-sunflower cropping system. Indian Journal of Weed Science 30(3 and 4), 103–107.
- Rekha, K.B., Raju, M.S., Reddy, M.D., 2002. Effect of herbicides in transplanted rice. Indian Journal of Weed Science 34(1 and 2), 123–125.
- Rout, D., Satapathy, 1996. Chemical weed control in rainfed maize (*Zea mays* L.). Indian Journal of Agronomy 41(1), 51–53.
- Satorre, E.H., de la Fuente, E.B., Mas, M.T., Suarez, S.A., Kruk, B.C., Guglielmini, A., Verdu, A.M.C., 2020. Crop rotation effects on weed communities of soybean (*Glycine max* L. Merr.) agricultural fields of the Flat Inland Pampa. Crop Protect. 130, 105068.
- Saha, S., Rao, K.S., 2009. Efficacy of sulfonylurea herbicides for broad-spectrum weed control in wet direct-sown summer rice. Oryza-An International Journal on Rice 46(2), 116–119.
- Saikia, T.P., Pandey, J., 2001. Annals of Agricultural Research New Series 22(2), 262–266.
- Saini, J.P., Punam, R., Chadha, S., Sharma, S., Bhardwaj, N., Rana, N., 2013. Non-chemical methods of weed management in maize under organic production system. Indian Journal of Weed Science 45(3) 198–200.
- Sasikala, K., Boopathi, S.N.M., Ashok, P., 2014. Evaluation of methods of sowing and post emergence herbicides for efficient weed control in zero till sown rice fallow black gram *Vigna mungo* L. International Journal of Farm Sciences 4(1), 81–91.
- Sharma, S.D., Punia, S.S., Malik, R.K., Narwal, S., 2004. Efficacy of cyhalofop-butyl against weeds in rice nursery. Indian Journal of Weed Science 36, 181–183.
- Sharma, A.R., Toor, A.S., Sur, H.S., 2000. Effect of interculture operations and scheduling of atrazine application on weed control and productivity of maize (*Zea mays*) in Shiwalik foothills of Punjab. Indian Journal of Agricultural Sciences 70, 757–761.
- Shekhar, J., Mankotia, B.S., Bindra, A.D., 2004. Bio-efficacy of some new herbicides against weed in transplanted rice (*Oryza sativa* L.). Indian Journal of Weed Science 36(1 and 2), 50–53.
- Sheoran, P., Sardana, V., Kumar A., Mann, A., Singh, S., 2015. Integrating herbicidal and conventional approach for profitable weed management in groundnut (*Arachis hypogaea*). Indian Journal of Agronomy 60(4), 110–113.
- Singh, G., Kaur, H., Khanna, V., 2014. Weed management in lentil with post-emergence herbicides. Indian Journal of Weed Science 46(2), 187–189.
- Singh, Y., Singh, V.P., Singh, G., Yadav, D.S., Sinha, R.K.P., Johnson, D.E., Mortimer, A.M., 2011. The

- implications of land preparation, crop establishment method and weed management on rice yield variation in the rice-wheat system in the Indo-Gangetic plains. *Field Crops Research* 121(1), 64–74.
- Singh, G., Singh, O.P., Singh, S., Prasad, K., 2010. Weed management in late sown wheat (*Triticum aestivum*) after rice (*Oryza sativa*) in rice-wheat system in rainfed lowland. *Indian Journal of Agronomy* 55(2), 83–88.
- Singh, R.K., Singh, R.P., Mukherjee, D., 2009. Effect of fertilizer placement and weed management practice on weed dynamics and yield of rainfed mustard (*Brassica juncea* L.). *Environment and Ecology* 27(4B), 1932–1938.
- Singh, R., 2006. Use of soil solarization in weed management on soybean under Indian conditions. *Tropical Science* 46(2), 70–73.
- Singh, S., Singh, G., Singh, V.P., Singh, A.P., 2005. Effect of establishment methods and weed management practices on weeds and rice in rice-wheat cropping system. *Indian Journal of Weed Science* 37, 51–57.
- Singh, G., Deka, J., Singh, D., 1988. Response of upland rice to seed rate and butachlor. *Indian Journal of Weed Science* 20(1), 23–30.
- Srinivasan, G., Subbian, P., 1991. Study on weed management in rice pulse cropping system. *Journal of Maharashtra Agricultural Universities* 16(3), 376–378.
- Subramanian, E., Martin, G., Balasubramanian, R., 2006. Effect of integrated weed management practices on growth and their residual effect on succeeding pulse crop. *Indian Journal of Agronomy* 51(2), 93–96.
- Subramanyam, D., Reddy, C.R., Reddy, D.S., 2007. Influence of puddling intensity and water management practices on weed dynamics and yield of transplanted rice (*Oryza sativa* L.). *Indian Journal of Agronomy* 2(3), 225–230.
- Talnikar, A.S., Kadam, G.L., Karande, D.R., Jogdand, P.B., 2008. Integrated weed management in pigeonpea [*Cajanus cajan* (L.) Millsp.]. *International Journal of Agricultural Sciences* 4(1), 363–370.
- Thiyagarajan, T.M., Kumar, S.K., Bindraban, P.S., Hengsdijk, S., Ramasamy, V., Velu, D., Durgadevi, K., Govindarajan, R., Priyadarshini, R., Sudhalakshmi, C., Nisha, P.T., Gayathry, G., 2002. Crop management options for increasing water productivity in rice. *Journal of Agriculture Research and Management* 1(4), 169–181.
- Walia, U.S., Brar, L.S., 2006. Effect of tillage and weed management on seed bank of *Phalaris minor* Retz. in wheat under rice-wheat sequence. *Indian Journal of Weed Science* 38(1 and 2), 104–107.
- Yadav, R.B., Singh, R.V., Yadav, K.G., 2013. Weed management in lentil. *Indian Journal of Weed Science* 45(2), 113–115.
- Yaduraju, N.T., 2006. Herbicide resistant crops in weed management. In: The extended summaries, golden jubilee national symposium on conservation agriculture and environment. October 26–28, Banaras Hindu University, Banaras, 297–98.
- Yaduraju, N.T., Kulshrestha, G., Mani, V.S., 1980. Herbicide studies in groundnut. *Indian Journal of Agronomy* 25, 447–52.