



Response of Foliar Nutrient Management on Yield, Nutrient Uptake and Use Efficiency of *Utera* Rapeseed-Mustard Varieties Grown in Rice–Fallows of Lower Indo-Gangetic Plains

Somnath Sah^{ID}, Arun Kumar Barik[✉], Siddharth Sankar Patre

Dept. of Agronomy, Palli Siksha Bhavana (Institute of Agriculture), Visva- Bharati, Sriniketan, Birbhum, West Bengal (731 236), India



Corresponding ✉ arunkumar.barik@visva-bharati.ac.in

^{ID} 0000-0002-0840-8858

ABSTRACT

A two-year field experiment during winter seasons (November to February) of 2020–2021 and 2021–2022 was conducted to evaluate the effect of foliar spray of macro and micro nutrients on different rapeseed-mustard varieties under relay cropping in rice fallows. The results revealed that foliar spray of macro and micro nutrients had significant effect on seed yield, oil content, oil yield, nutrient uptake and use efficiency irrespective of rapeseed mustard varieties. Variety *Binoy* (B-9) registered significantly higher oil content (38.1%) over other two varieties but PM 28 showed its superiority when total oil yield per unit area was considered. Mustard variety PM 28 exhibited significantly higher total nitrogen (48.78 kg ha⁻¹), phosphorus (11.79 kg ha⁻¹), potassium (45.97 kg ha⁻¹) and sulphur (23.29 kg ha⁻¹) uptake both in seeds as well as in stover. Combined application of macro and micro nutrients *i.e.*, spraying of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS exhibited significantly higher seed yield, oil content, protein content, nutrient uptake of N, P, K, S, Zn and nitrogen use efficiency than other foliar spray of nutrients. Rapeseed-mustard variety PM-28 along with foliar application of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS performed best in rice fallow of lower Indo-Gangetic plains.

KEYWORDS: NUE, nutrient uptake, oil yield, spray nutrients

Citation (VANCOUVER): Saha et al., Response of Foliar Nutrient Management on Yield, Nutrient Uptake and Use Efficiency of *Utera* Rapeseed-Mustard Varieties Grown in Rice–Fallows of Lower Indo-Gangetic Plains. *International Journal of Bio-resource and Stress Management*, 2025; 16(1), 01-08. [HTTPS://DOI.ORG/10.23910/1.2025.5665](https://doi.org/10.23910/1.2025.5665).

Copyright: © 2025 Saha et al. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, that permits unrestricted use, distribution and reproduction in any medium after the author(s) and source are credited.

Data Availability Statement: Legal restrictions are imposed on the public sharing of raw data. However, authors have full right to transfer or share the data in raw form upon request subject to either meeting the conditions of the original consents and the original research study. Further, access of data needs to meet whether the user complies with the ethical and legal obligations as data controllers to allow for secondary use of the data outside of the original study.

Conflict of interests: The authors have declared that no conflict of interest exists.

RECEIVED on 01st August 2024 RECEIVED in revised form on 15th December 2024 ACCEPTED in final form on 02nd January 2025 PUBLISHED on 16th January 2025

1. INTRODUCTION

The Indo-Gangetic plain (IGP) of India, spread over 44 million ha, is the food basket of South Asia. Approximately 76% of its area falls in India, dominated by Punjab, Haryana, Uttar Pradesh, Bihar and West Bengal (Kumar et al., 2021). The IGP regions of eastern India comprises alluvial soil, predominant with rice-based cropping systems which are prevailing years after years with locally available (Midya et al., 2021) long duration rice cultivars. This triggers delayed sowing of subsequent crops resulting poor yield. A sustainable production is needed to balance world's nutritional needs while reducing agriculture's negative environmental impact (Liu et al., 2022). Adopting conservation or sustainable farming, *utera* rapeseed-mustard can easily be raised effectively and efficiently just after *aman* paddy. It might have emphasized to adapt in large scale for increasing cropping intensity in Eastern India (Kumar et al., 2019). In spite of advantageous role of tillage, conservation agriculture is getting impetus of having benefits of undisturbed soil (Ramesh et al., 2021). *Utera* cropping minimizes tillage cost and taking advantages of residual fertility and moisture (Venkateswarlu et al., 2009). Crop varieties efficient in extracting nutrients from soil for growth are the cornerstone for future global crop production (Lyzenga et al., 2023). Apart from sowing time, varieties plays a crucial role to realize potential yield of rapeseed-mustard (Sharma and Kumar, 2023, Saha et al., 2023). Adopting smart agronomic tools like foliar spray of nutrients also augment the yield of *utera* rapeseed-mustard (Saha et al., 2023).

Rapeseed-mustard is the leading edible oilseed crop (Kumar et al., 2014) after groundnut, contributing nearly 26.1% of total oilseed production in India (Meena et al., 2018 and Meena et al., 2013). Eight different species are being cultivated all over the globe (Singh et al., 2023). The crop responded well to applied major plant nutrients (N, P and K) along with sulfur, zinc and boron as they plays vital roles in production phenology of crops and show positive response on appliance of sulfur, zinc and boron (Dhaliwal et al., 2021, Karthika et al., 2020, Pati and Mahapatra, 2015). Foliar fertilization is more effective, economical (Fageria et al., 2009) and successful in improving yields by delaying senescence and converting late flushes of flower into economic parts (Banerjee et al., 2019). It can make nutrients available directly to the site of food synthesis with minimum wastage (Saha et al., 2023). Foliar feeding of micronutrients is cheap, more effective, efficient and having lesser impact to pollution as it adds less fertilizers to the environment (Bhatt et al., 2020 and Aziz et al., 2019). Foliar application of nitrogenous fertilizers would not only reduce soil pollution but also improves uptake of nitrogen

(Sunkad et al., 2023). Rapeseed-mustard is very sensitive to micronutrient deficiency, especially zinc and boron (Rathore et al., 2015). S (sulfur), Zn, B and Fe are important plant nutrients to optimize productivity of rapeseed-mustard (Meena et al., 2022b). The oil content of rapeseed-mustard is being largely governed by mineral fertilization of the crops (Olama et al., 2014). On the other hand, NPK fertilizers along with micronutrients zinc (Zn) and boron (B) contributes an important role in nutrient uptake and nutrient use efficiency in oilseed crops (Immanuel, 2019). K application can improve nutrient use efficiency (NUE) of other nutrients with a capacity to mitigate adverse effects of excessive N on disease and insect-pest incidences, improving crop yields, crop health and NUE (Perelman et al., 2021).

In this experiment, an attempt was made to study the effect of foliar spray of nutrients on different varieties of rapeseed-mustard with reference to nutrient content and uptake, seed oil content and protein content of the crop.

2. MATERIALS AND METHODS

A field experiment was conducted during winter seasons (November to February) of 2020–21 and 2021–22 at Pulses and Oilseeds Research Station, Berhampore, West Bengal 742101, India with 24° 6' N latitude and 88° 15' E longitude and average altitude of 41 m above mean sea level. The experiment was designed in a splitplot design keeping rapeseed-mustard varieties, viz., $V_1 = \text{Anushka}$, $V_2 = \text{Biney}$ (B-9) and $V_3 = \text{PM-28}$ in main plots and eight different foliar spray of nutrients viz. F_0 : No foliar spray, F_1 : Foliar spray with N:P:K (19:19:19) @ 2% at 30 DAS and 45 DAS, F_2 : Foliar spray with ZnSO_4 @ 0.5% at 40 DAS, F_3 : Foliar spray with Boron (20%) @ 0.1% at 50 DAS, F_4 : $F_1 + F_2$, F_5 : $F_1 + F_3$, F_6 : $F_2 + F_3$, F_7 : $F_1 + F_2 + F_3$ in sub-plots (Table 1). Each main plots and sub-plots were replicated thrice in the experimental field having plots of size 5×4 m². The weather of the experimental site was under sub-tropical climate having average annual rainfall ranging from 1200 to 1300 mm concentrating during rainy months from June to September. During the crop growth period the mean maximum temperature varied between 21.3°C to 35.4°C, and minimum between 10.7°C to 26.5°C. The maximum and minimum relative humidity ranged between 71.4% to 95.4% and 38.6% to 80.1%, respectively.

The soil of the experimental site was silty clay loam in texture, neutral in reaction (pH 7.46), having organic carbon, available nitrogen, phosphate and potash with 0.40%, 299 kg ha⁻¹, 45 kg ha⁻¹ and 278 kg ha⁻¹ respectively.

The seeds of different varieties of rapeseed-mustard were sown in the standing rice field 10 days before harvesting of rice. Nitrogen, phosphorus and potassium were applied in the form of straight fertilizers i.e. urea,

Table 1: Effect of different varieties and foliar spray of nutrients on seed yield, oil content (%), oil yield (kg ha⁻¹), protein content (%), protein harvest (kg ha⁻¹), PFP (kg kg⁻¹), AE (kg kg⁻¹) and ARE of relay rapeseed-mustard

Treatments	Seed yield (kg ha ⁻¹)	Oil content (%)	Oil yield (kg ha ⁻¹)	Protein content (%)	Protein harvest (kg ha ⁻¹)	PFP (kg kg ⁻¹)	AE (kg kg ⁻¹)	ARE
Rapeseed-mustard varieties (V)								
V ₁	817.98	37.1	305.5	13.03	108.3	15.59	3.41	0.20
V ₂	960.46	38.1	368.2	13.67	132.2	18.29	5.23	0.27
V ₃	1198.46	33.5	404.1	16.22	194.7	22.87	5.17	0.27
SEm±	13.09	0.17	3.9	0.07	2.19	0.242	0.31	0.012
CD (p=0.05)	42.70	0.55	12.8	0.27	7.14	0.79	1.00	0.040
Foliar spray of nutrients (F)								
F ₀	774.22	31.9	245.0	11.70	93.1	15.48	-	-
F ₁	1012.94	35.4	355.2	14.20	145.4	18.57	5.00	0.22
F ₂	872.00	36.0	310.1	13.89	121.3	17.44	2.86	0.15
F ₃	891.50	35.5	312.3	13.65	122.0	17.83	2.90	0.16
F ₄	1076.72	38.0	406.3	15.70	171.4	19.74	6.06	0.34
F ₅	1130.06	37.9	425.8	15.11	172.3	20.71	6.84	0.37
F ₆	947.72	36.2	340.7	13.83	132.7	18.95	4.23	0.21
F ₇	1233.22	38.9	479.0	16.36	202.4	22.60	8.93	0.51
SEm±	25.83	0.43	10.1	0.09	4.11	0.494	0.44	0.017
CD (p=0.05)	72.64	1.20	28.3	0.25	11.57	1.389	1.23	0.047
Interaction V×F								
SEm±	44.74	0.738	17.4	0.19	7.126	0.855	0.76	0.029
CD (p=0.05)	NS	NS	NS	NS	NS	NS	NS	NS
Interaction F×V								
SEm±	43.85	0.711	16.8	0.15	7.016	NS	NS	NS
CD (p=0.05)	NS	NS	NS	NS	NS			

single super phosphate and muriate of potash at 50% of the recommended dose (100:50:50 kg ha⁻¹) just five days before sowing of rapeseed mustard seeds when soil was saturated with excess moisture. The foliar application was done according to the treatments. Irrigation to each plot was uniformly applied as life savers. Plant protection measures were taken as and when required and seeds were treated with fungicides irrespective of treatments.

The seed samples from each treatment were taken for oil estimation with the help of Soxhlet's apparatus (Anonymous, 1970). The nutrient content of seed and stover (N, P, K, S and Zn content) were estimated in both the years and uptake was calculated with respective nutrient content (Singh et al., 2005). Protein content of the grain was driven out by multiplying the nitrogen content with the multiplication factor 6.25. Further the protein harvest per unit area (kg ha⁻¹) was calculated by multiplying the protein content with seed yield. Partial factor productivity

(PFP), agronomic efficiency (AE) and apparent recovery efficiency (ARE) of applied nutrients were calculated using the following formulae suggested by Baligar et al. (2001), Dobermann (2007), Fageria et al. (2008) and Fageria et al. (2015). Standard statistical methods of analysis were followed (Panse and Sukhatme, 1978; Gomez and Gomez, 1984).

3. RESULTS AND DISCUSSION

3.1. Seed yield

A significant difference was found among tested varieties of rapeseed-mustard on seed yield (Table 1). Among three tested varieties the highest seed yield was achieved by PM-28 (1198.46 kg ha⁻¹) which was significantly higher than others. Higher seed yield achieved by PM-28 might be attributed to higher growth and yield attributes, longer duration and higher nutrient uptake than other two varieties. Foliar spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and

45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS revealed significantly highest seed yield across varieties. Combined application of macro and micro resulted higher seed yield might be due to the synergistic effect. Aziz et al. (2019) also found similar findings of increased tillering, spike length, grain yield in wheat. Foliar fertilizers used in the winter oilseed rape affected an increase in yield on average ranging from 2.5% to 18.5% (Sikorska et al., 2020). Ashkiani et al. (2020) found that application ZnSO₄ had a positive effect on rapeseed-mustard seed yield and oil yield. Meena et al. (2022b) confirmed that application of secondary (S) and micronutrients (Zn and B) significantly enhanced yield (28.5 q ha⁻¹) in rapeseed-mustard.

3.2. Oil content and oil yield

Among various test varieties different foliar spray of nutrients significantly influenced the oil content in seed as well as the oil yield irrespective of all three tested varieties. Highest oil content was found in B-9 (38.1%) followed by *Anushka* (37.1%) and PM-28 (33.5%). Among different combinations of foliar spray of nutrients spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS showed significantly higher seed oil content (38.9%). The trend was similar in all the three tested varieties (Table 1). Considering the total oil yield mustard variety PM-28 (404.1 kg ha⁻¹) out yielded other two yellow sarson varieties viz. B-9 (368.2 kg ha⁻¹) and *Anushka* (305.5 kg ha⁻¹). Among the different foliar spray of nutrients spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS proved its superiority over others (Table 1). Cakmak (2008) found that application of Zn had a significant effect on seed oil content. The application of Zn had a significant effect on rapeseed-mustard oil and oil content of seeds increased with Zn application (Ahmadi, 2010). Application of foliar nutrients might have positive effect for enhancing metabolic and photosynthetic activity due to their involvement in chlorophyll formation, different enzyme activation as well as primary and secondary metabolism, and thus, ultimately increased the oil quantity (Rana et al., 2020). Their findings aligned with findings of Sarkar et al. (2021) were they found that foliar nutrient spray significantly increase the oil percentage (39.9%) in hybrid mustard (Kesari-5111) and lowest oil content (33.7%) was obtained with no foliar spray treatment. Meena et al. (2022b) observed positive effect of nutrient enrichment of FYM with micro (Zn+B) and secondary (S) nutrients for maximizing oil content (43.1%) and seed yield in rapeseed-mustard.

3.3. Protein content and protein yield

The protein content in seed and protein harvest by crop were significantly influenced due to different foliar spray of nutrients and the trends were similar in all the three

tested varieties. Highest protein content was found in PM-28 (16.22%) followed by B-9 (13.67%) and *Anushka* (13.03%). Among different combinations of foliar spray of nutrients spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS showed significantly higher seed protein content (16.36%). The trend was similar in all the three tested varieties. Considering the total protein yield mustard variety PM-28 (194.7 kg ha⁻¹) gave significantly higher protein yield than other two yellow sarson varieties viz. B-9 (132.2 kg ha⁻¹) and *Anushka* (108.3 kg ha⁻¹). Among the different foliar spray of nutrients spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS (202.4 kg ha⁻¹) proved its superiority over others (Table 1). The positive effect of balanced nutrients supply of NPK along with S and Zn on seed and quality yield were also reported by Tatarwal et al. (2013). Zinc is directly or indirectly required by several enzyme systems and is closely involved in the nitrogen metabolism which ultimately enhanced the synthesis of many enzymes and proteins in plant body (Mondal et al., 2020). Meena et al. (2022a) in an experiment when they applied 125% RDF of NPK along with S and Zn got higher seed protein yield in *Brassica* (Indian mustard). Conversely, the combined effect of macro and micro nutrients played a very significant role by partitioning of photosynthates due to their synergistic effect (Singh et al., 2020) which also helped in better yielding of seed protein.

3.4. Nutrient uptake

From two years of field experiments a significant difference in nutrient uptake by seeds, stover and total nutrient uptake by rapeseed-mustard were noticed among the tested varieties (Figure 1 to 5). Mustard variety PM 28 showed significantly higher total nitrogen (48.78 kg ha⁻¹), phosphorus (11.79 kg ha⁻¹), potassium (45.97 kg ha⁻¹) and sulphur (23.29 kg ha⁻¹) uptake when compared with other two varieties. Although total zinc uptake was significantly higher in B-9 (116.9 g ha⁻¹). Among different combinations of foliar spray of nutrients, foliar spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS exhibited significantly higher total nutrient uptake than other foliar spray of nutrients in different varieties of rapeseed-mustard. No significant interaction was noticed in nutrient uptake of *relay* rapeseed mustard in rice fallows. Ganie et al. (2014) opined that boron nutrition significantly improved the N, P, K, S and B uptake in French bean. Chaudhary et al. (2016) and Meena et al. (2022a) reported, foliar application of nutrients ensured balanced and continuous supply of essential plant nutrients throughout the growth periods. This attributed higher






Figure 1: Total N uptake (kg ha⁻¹) by *utera* rapeseed-mustardFigure 2: Total P uptake (kg ha⁻¹) by *utera* rapeseed-mustardFigure 3: Total K uptake (kg ha⁻¹) by *utera* rapeseed-mustard

Figure 4: Total S uptake (kg ha⁻¹) by *utera* rapeseed-mustard uptake of nutrients like N, P, K, S by rapeseed-mustard. Higher accumulation and uptake of nutrients under these treatments could be ascribed to better availability and synergistic effect of applied nutrients (Kumaran and Solaimalai, 2000; Samant and Panigrahi, 2018). Marked enhancement in N, P and K uptake in rapeseed-mustard was due to application of sulfur and boron which attributed to better utilization of nutrients through the

Figure 5: Total Zn uptake (g ha⁻¹) by *utera* rapeseed-mustard

well-developed root system with a concomitant increase in higher yield and higher concentration of N, P and K in both seed and stover (Mathew and George, 2013; Jaiswal et al., 2015; Basumatary et al., 2021).

3.5. Nitrogen use efficiency (NUE)

Among different tested varieties of rapeseed-mustard, PM-28 revealed significantly higher partial factor productivity (22.87 kg kg⁻¹). While assessing the agronomic efficiency, B-9 showed significantly higher value (5.23 kg kg⁻¹) than *Anushka* (3.41 kg kg⁻¹) but at par with PM-28 (5.17 kg kg⁻¹). B-9 and PM-28 showed at par values of Apparent Recovery Efficiency (0.27) but significantly higher than *Anushka* (0.20). Partial Factor Productivity (PFP), Agronomic efficiency (AE) and Apparent Recovery Efficiency (ARE) were significantly influenced by application of different combinations of foliar spray of nutrients significantly higher values were exhibited by the treatment having foliar spray of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS over rest of the treatments on pooled data basis (22.60 kg kg⁻¹ 8.93 kg kg⁻¹ and 0.51, respectively, Table 1). High nutrient efficient cultivars facilitated a production system with sustainability as reported by Adhikari et al. (2023). By applying foliar spray of essential micronutrients like Zn and B along with NPK fertilizers, Immanuel (2019) succeeded to boost and sustain seed yield of sunflower through increasing nutrient uptake and nutrient use efficiency. Many long-term field trials have proven how K application can also boost NUE of P and other nutrients like S as reported by Perelman et al. (2021). Their studies had also shown that K can lessen negative effects of excess N applied to crop, thereby enhancing yields and health of crops ultimately the NUE of N.

4. CONCLUSION

Rapeseed-mustard variety PM-28 with combined application of N:P:K (19:19:19) @ 2% twice at 30 DAS and 45 DAS+ZnSO₄ @ 0.05% at 40 DAS+Boron (20%) @ 0.1% at 50 DAS performed best for obtaining higher seed yield, oil yield, protein yield, nutrient uptake and use efficiency over other varieties and enhancing productivity

crop quality of *utera* rapeseed mustard in rice fallows of lower Indo-Gangetic Plains.

5. REFERENCES

- Adhikari, S., Anuragi, H., Chandra, K., Tarte, S.H., Dhaka, S.R., Jatav, H.S., Hingonia, K., 2023. Molecular basis of plant nutrient use efficiency-concepts and challenges for its improvement-chapter six. In: Aftab, T., Hakeem, K.R. (Eds.), Sustainable plant nutrition. Academic Press, 107–151, ISBN 9780443186752. <https://doi.org/10.1016/B978-0-443-18675-2.00001-8>.
- Ahmadi, M., 2010. Effect of zinc and nitrogen fertilizer rates on yield and yield components of oilseed rape (*Brassica napus* L.). American-Eurasian Journal of Agricultural & Environmental Sciences 7(3), 259–264. [https://www.idosi.org/aejaes/jaes7\(3\)/3.pdf](https://www.idosi.org/aejaes/jaes7(3)/3.pdf).
- Anonymous, 1970. Official methods of analysis of the association of official analytical chemists. 11th ed. Editor: W. Horwitz. Washington D.C., Pub: Association of Official Analytical Chemists. 1970. Available from <https://doi.org/10.1002/jps.2600600253> Accessed on 24.07.2024.
- Ashkiani, A., Sayfzadeh, S., Rad, A.H.S., Valadabadi, A., Masouleh, E.H., 2020. Effects of foliar zinc application on yield and oil quality of rapeseed genotypes under drought stress. Journal of Plant Nutrition 43(11), 1594–1603. DOI: 10.1080/01904167.2020.1739299.
- Aziz, M.Z., Yaseen, M., Abbas, T., Naveed, M., Mustafa, A., Hamid, Y., Saeed, Q., Ming-gang, X.U., 2019. Foliar application of micronutrients enhances crop stand, yield and the biofortification essential for human health of different wheat cultivars. Journal of Integrative Agriculture 18(6), 1369–1378.
- Baligar, V.C., Fageria, N.K., He, Z.L., 2001. Nutrient use efficiency in plants. Communications in Soil Science and Plant Analysis 32(7–8), 921–950.
- Banerjee, P., Visha Kumari, V., Nath, R., Bandyopadhyay, P., 2019. Seed priming and foliar nutrition studies on relay grass pea after winter rice in lower Gangetic plain. Journal of Crop and Weed 15(3), 72–78.
- Basumatary, A., Chauhan, S., Bhupenchandra, I., Das, K.N., Ozah, D.J., 2021. Impact of sulfur and boron fertilization on yield, quality of crop and nutrient use efficiencies in rapeseed in subtropical acidic soil of Assam, India. Journal of Plant Nutrition 44(12), 1779–1793. DOI: 10.1080/01904167.2021.1884708.
- Bhatt, R., Hossain, A., Sharma, P., 2020. Zinc bio-fortification as an innovative technology to alleviate the zinc deficiency in human health: a review. Open Agriculture 5(1), 176–187.
- Cakmak, I., 2008. Enrichment of cereal grains with zinc: agronomic or genetic bio-fortification. Plant and Soil 302, 1–17.
- Chaudhary, S., Shukla, A., Bhushan, C., Negi, M.S., 2016. Assessment of the system of root intensification in rapeseedmustard (*Brassica juncea*). Indian Journal of Agronomy 61(1), 119–122.
- Dhaliwal, S.S., Sharma, V., Shukla, A.K., Verma, V., Behera, S.K., Sandhu, P.S., Kaur, K., Gaber, A., Althobaiti, Y.S., Abdelhadi, A.A., Hossain, A., 2021. Assessment of agro-economic indicators of *Sesamum indicum* L. as influenced by application of boron at different levels and plant growth stages. Molecules 26(21), 6699. <https://doi.org/10.3390/molecules26216699>.
- Dobermann, A., 2007. Nutrient use efficiency-measurement and management. In: Krauss, A., Isherwood, K., Heffer, P. (Eds.), Fertilizer best management practices: general principles, strategy for their adoption and voluntary initiatives versus regulations. International Fertilizer Industry Association, Paris, France, 1–28.
- Fageria, N.K., Baligar, V.C., Heinemann, A.B., Carvalho, M.C.S., 2015. Nitrogen uptake and use efficiency in rice. In: Rakshit, A., Singh, H.B., Sen, A. (Eds.), Nutrient use efficiency: from basics to advances. Springer, New Delhi, 285–296. https://doi.org/10.1007/978-81-322-2169-2_18.
- Fageria, N.K., Baligar, V.C., Li, Y.C., 2008. The role of nutrient efficient plants in improving crop yields in the twenty first century. Journal of Plant Nutrition 31(6), 1121–1157.
- Fageria, N.K., Barbosa Filho, M.P., Moreira, A., Guimaraes, M., 2009. Foliar fertilization of crops. Journal of Plant Nutrition 32(6), 1044–1064.
- Ganie, M.A., Akhter, F., Bhat, M.A., Najar, G.R., 2014. Growth yield and quality of french bean (*Phaseolus vulgaris* L.) as influenced by sulphur and boron application on inceptisols of Kashmir. The Bioscan 9(2), 513–518.
- Gomez, K.A., Gomez, A.A., 1984. Statistical procedures for agricultural research (2nd Edn.), An International Rice Research Institute Book, a Wiley-inter-science Publication, John Wiley & Sons. New York. Chichester. Brisbane. Toronto. Singapore. February 1984, ISBN: 978-0-471-87092-0, 704.
- Immanuel, R.R., 2019. Nutrient uptake and nutrient use efficiency of sunflower in response to Zn and B micronutrient fertilization. Plant Archives 19(2), 543–547.
- Jaiswal, A.D., Singh, S.K., Singh, Y.K., Singh, S., Yadav, S.N., 2015. Effect of sulphur and boron on yield and quality of mustard (*Brassica juncea* L.) grown on vindhyan red soil. Journal of the Indian Society of Soil Science 63(3), 362–364. doi: 10.5958/0974-0228.2015.00047.X. Available from <https://www>.

- indianjournals.com/ijor.aspx?target=ijor:jisss&volume=63&issue=3&article=014&type=pdf
- Karthika, K.S., Philip, P.S., Neenu, S., 2020. Brassicaceae plants response and tolerance to nutrient deficiencies. In: Hasanuzzaman, M. (Ed.), The plant family brassicaceae: biology and physiological responses to environmental stresses. Springer Nature Singapore Pte Ltd., 337–362. https://doi.org/10.1007/978-981-15-6345-4_11.
- Kumar, N., Chhokar, R.S., Meena, R.P., Kharub, A.S., Gill, S.C., Tripathi, S.C., Gupta, O.P., Mangrauthia, S.K., Sundaram, R.M., Sawant, C.P., Ajita Gupta, A., Naorem, A., Kumar, M., Singh, G.P., 2021. Challenges and opportunities in productivity and sustainability of rice cultivation system: a critical review in Indian perspective. Cereal Research Communication 50, 573–601. doi: 10.1007/s42976-021-00214-5.
- Kumar, P., Kumar, A., Kumar, S., Kumar, P., 2014. Effect of zinc and iron application on yield and acquisition of nutrient on mustard crop (*Brassica juncea* L.). Journal of Plant Development Sciences 6(3), 413–416.
- Kumar, R., Mishra, J.S., Rao, K.K., Bhatt, B.P., Hazra, K., Hans, H., Mondal, S., 2019. Sustainable intensification of rice fallows of Eastern India with suitable winter crop and appropriate crop establishment technique. Environmental Science and Pollution Research 26, 29409–29423. <https://doi.org/10.1007/s11356-019-06063-4>.
- Kumaran, S., Solaimalai, A., 2000. Effect of organic manure and inorganic fertilizers on yield and nutrient uptake of irrigated groundnut. Crop Research (Hisar) 20(1), 35–38.
- Liu, Q., Wu, K., Song, W., Zhong, N., Wu, Y., Fu, X., 2022. Improving crop nitrogen use efficiency toward sustainable green revolution. Annual Review of Plant Biology 73, 523–551. <https://doi.org/10.1146/annurev-arplant-070121-015752>.
- Lyzenga, W.J., Liu, Z., Olukayode, T., Zhao, Y., Kochian, L.V., Ham, B.K., 2023. Getting to the roots of N, P, and K uptake. Journal of Experimental Botany 74(6), 1784–1805. <https://doi.org/10.1093/jxb/erad035>.
- Mathew, J., George, S., 2013. Synergistic-influence of sulphur and boron on enhancing the productivity of sesame (*Sesamum indicum*) grown in an Entisol of Kerala. Journal of the Indian Society of Soil Science 61(2), 122–127.
- Meena, B.S., Narolia, R.S., Nagar, G., Meena, S.N., Meena, D.S., 2022a. Effect of sowing window, planting geometry and nutrient-management on yield, nutrient uptake and economics of Indian mustard in Vertisols of south-eastern Rajasthan. Indian Journal of Agronomy 67(1), 50–57.
- Meena, D.S., Meena, V.R., Meena, A.K., 2013. Fertilizer management studies on growth and productivity of hybrid Indian mustard *Brassica juncea* (L.). Journal of Oilseed Brassica 4(1), 39–42.
- Meena, M., David, A.A., Kumar, S., 2018. Effect of different levels of NPK and zinc sulphate on yield and oil content in mustard (*Brassica juncea* L.) Var. Jai Kisan. International Journal of Pure and Applied Biosciences 6(6), 722–727.
- Meena, V.S., Meena, M.K., Dotaniya, M.L., Meena, M.D., Jat, R.S., Choudhary, R.L., Singh, H., Meena, V.D., Rai, P.K., 2022b Profitability and productivity of Indian mustard affected by micro and secondary nutrient application in calcareous soil. Journal of Oilseed Brassica 13(1), 53–58. Available from <https://krishi.icar.gov.in/jspui/bitstream/123456789/76487/1/14.%20Profitability%20and%20productivity%20of%20Indian%20mustard%20affected%20by%20micro%20and%20secondary%20nutrient%20application%20in%20calcareous%20soil.pdf>.
- Midya, A., Saren, B.K., Dey, J.K., Maitra, S., Praharaj, S., Gaikwad, D.J., Gaber, A., Alhomrani, M., Hossain, A., 2021. Crop establishment methods and integrated nutrient management improve: Part II. Nutrient uptake and use efficiency and soil health in rice (*Oryza sativa* L.) field in the lower Indo-Gangetic Plain, India. Agronomy 11(9), 1894. <https://doi.org/10.3390/agronomy11091894>
- Mondal, R., Goswami, S., Goswami, S.B., Jana, K., 2020. Effect of different nutrient management practices on growth, grain yield, production economics, soil nutrient availability of transplanted *kharif* rice (*Oryza sativa* L.) and correlation studies. Journal of Crop and Weed 16(1), 172–179.
- Olama, V., Ronaghi, A., Karimian, N., Yasrebi, J., Hamidi, R., Tavajjoh, M., Kazemi, M.R., 2014. Seed quality and micronutrient contents and translocations in rapeseed (*Brassica napus* L.) as affected by nitrogen and zinc fertilizers. Archives of Agronomy and Soil Science 60(3), 423–435.
- Panase, V.G., Sukhatme, P.V., 1978. Statistical methods for agricultural workers (3rd Edn.). Indian Council of Agricultural Research.
- Pati, P., Mahapatra, P.K., 2015. Yield performance and nutrient uptake of Indian mustard (*Brassica juncea* L.) as influenced by integrated nutrient management. Journal of Crop and Weed 11(1), 58–61.
- Perelman, A., Imas, P., Bansal, S.K., 2021. Role of potassium for improving nutrient use efficiency in agriculture. In: Bhatt, R., Meena, R.S., Hossain, A., (Eds.), Input use efficiency for food and environmental security. Springer, Singapore. <https://doi.org/10.1007/978->

- 981-16-5199-1_13.
- Ramesh, K., Jayaraman, S., Bhaskar, S., 2021. Inbuilt mechanisms for managing weeds in conservation agriculture systems: a revisit. In: Jayaraman, S., Dalal, R.C., Patra, A.K., Chaudhari, S.K. (Eds.), Conservation agriculture: a sustainable approach for soil health and food security. Springer, Singapore, 73–85. https://doi.org/10.1007/978-981-16-0827-8_4.
- Rana, K., Parihar, M., Singh, J.P., Singh, R.K., 2020. Effect of sulfur fertilization, varieties and irrigation scheduling on growth, yield, and heat utilization efficiency of Indian mustard (*Brassica juncea* L.). Communications in Soil Science and Plant Analysis 51(2), 265–275.
- Rathore, S.S., Shekhawat, K., Kandpal, B.K., Premi, O.P., 2015. Improving nutrient use efficiency in oilseeds brassica. In: Rakshit, A., Singh, H.B., Sen, A., (Eds.), Nutrient use efficiency: from basics to advances. Springer, New Delhi, 317–327.
- Ray, K., Sengupta, K., Pal, A.K., Banerjee, H., 2015. Effects of sulphur fertilization on yield, S uptake and quality of Indian mustard under varied irrigation regimes. Plant, Soil and Environment 61(1), 6–10.
- Saha, S., Barik, A.K., Kumar, B., 2023. Varietal performances and foliar application of nutrients on yield and economics of *Utera* rapeseed mustard in rice-fallow of alluvial soils. International Journal of Bio-resource and Stress Management 14(11), 1550–1555. <https://doi.org/10.23910/1.2023.4920>.
- Samant, T.K., Panigrahi, D., 2018. Effect of sowing dates and nutrient management practices on important parameters in Toria (*Brassica rapa* L. var. Toria). Journal of Oilseed Brassica 9(1), 22–26.
- Sarkar, A., Jana, K., Mondal, R., 2021. Growth and yield of hybrid mustard (*Brassica juncea* L.) as influenced by foliar nutrition in Gangetic plains of West Bengal. Journal of Crop and Weed 17(3), 35–40. <https://doi.org/10.22271/09746315.2021.v17.i3.1488>.
- Sharma, S.K., Kumar, A., 2023. Effect of date of sowing on growth, seed yield and economics of Indian mustard (*Brassica juncea*) varieties under rainfed Conditions. Indian Journal of Agricultural Research 57(1), 56–59. doi: 10.18805/IJAR.A-5947.
- Sikorska, A., Gugala, M., Zarzecka, K., 2020. Winter oilseed rape yield depending on foliar fertilization. Acta Scientiarum Polonorum Agricultura 19(1), 11–20. DOI: 10.37660/aspagr.2020.19.1.2.
- Singh, A.K., Singh, A., Aswin, C., Shashidhar, K.S., 2020. Evaluation of agro-chemicals for enhancing the productivity and profitability of late-sown Indian mustard (*Brassica juncea*) under acidic soils of Manipur. Indian Journal of Agronomy 65(1), 94–99.
- Singh, D., Chhonkar, P.K., Dwivedi, B.S., 2005. Manual on soil, plant and water analysis. Westville Publishing House, New Delhi, 1–200.
- Singh, M., Tomar, A., Kumar, S., 2023. Present status, production constraints and future research strategies in oilseed brassica. In: International Conference on Vegetable Oils 2023 (ICVO 2023)-Research, Trade, Value Chain and Policy January 17–21, 2023, Hyderabad. India Journal of Oilseeds Research 40 (Special Issue), 60–63. <https://doi.org/10.56739/jor.v40iSpecialissue.145143>
- Sunkad, P., Somanagouda, G., Channakeshava, R., Angadi, S.S., Banu, H., 2023. Effect and profitability of foliar application of thiourea on soybean yield. In: International Conference on Vegetable Oils 2023 (ICVO 2023)-Research, Trade, Value Chain and Policy January 17–21, 2023, Hyderabad, India. Pub. Journal of Oilseeds Research 40(Special Issue), 59–60.
- Tetarwal, J.P., Baldev, R., Meena, D.S., Tomar, S.S., 2013. Effect of moisture conservation and sulphur sources on productivity and water-use efficiency of Indian mustard (*Brassica juncea*) under rainfed conditions. Indian Journal of Agronomy 58(2), 231–236.
- Venkateswarlu, B., Sharma, K.L., Prasad, J.V.N.S., 2009. Conservation Agriculture—Constraints, Issues and Opportunities in Rainfed Areas, 4th World Congress on Conservation Agriculture—February, 4–7, 2009. New Delhi. Available from <http://www.irancon.org/wp-content/uploads/2021/01/4th-World-Congress-on-CA-India-4-7-Feb-2009.pdf>