



Effect of Supplementation of Onion Peel Powder as a Herbal Feed Additive on Carcass Traits, Sensory Evaluation and Cost Economics of Japanese Quails

Mujeebulla S. Md.¹, Susmita T.² , Tirupathi Reddy E.³ and Raja Kishore K.⁴

¹Dept. of Poultry Science, ²Dept. of Livestock Farm Complex, NTR CVSc, Gannavaram, Andhra Pradesh (521 102), India

³Dept. of Poultry Science, CVSc, Tirupati, Andhra Pradesh (517 502), India

⁴Dept. of Animal Nutrition, CVSc, Produtur, Andhra Pradesh (516 360), India



Corresponding  susmita.thullimalli@gmail.com

 0009-0001-0667-3025

ABSTRACT

The present study was carried out during November to December, 2023 at poultry unit, livestock farm complex, NTR College of Veterinary Science, Andhra Pradesh, to evaluate the effect of supplementation of onion peel powder (OPP) as an herbal feed additive on the performance of Japanese quail. 225 day-old Japanese quails were distributed randomly into five treatments with three replicates containing fifteen birds each and were fed with five experimental diets T₁ (Basal diet), T₂ (Basal diet+10 g kg⁻¹ OPP), T₃ (Basal diet+20 g kg⁻¹ OPP), T₄ (Basal diet+30 g kg⁻¹ OPP) and T₅ (Basal diet+40 g kg⁻¹ OPP) from day old to five weeks of age and were maintained under uniform management conditions. Two birds per replicate were slaughtered at the end of 5th week. The study showed that the carcass traits like pre-slaughter live weight, heart weight, and carcass weight significantly increased ($p < 0.01$) with increasing levels of OPP. The study found that supplementing OPP in diets from 0 to 40 g kg⁻¹ did not affect dressing percentage, liver, gizzard, giblet weights and sensory evaluation compared to other treatment groups. Quail fed with 40 g kg⁻¹ OPP had significantly lower feed cost kg⁻¹ gain. The study indicates that OPP can be safely supplemented up to 40 g kg⁻¹ in the diet without any negative impact on the performance of Japanese quail.

KEYWORDS: OPP, Japanese quails, carcass traits, dressing percentage, feed cost

Citation (VANCOUVER): Mujeebulla et al., Effect of Supplementation of Onion Peel Powder as a Herbal Feed Additive on Carcass Traits, Sensory Evaluation and Cost Economics of Japanese Quails. *International Journal of Bio-resource and Stress Management*, 2025; 16(3), 01-07. [HTTPS://DOI.ORG/10.23910/1.2025.5762](https://doi.org/10.23910/1.2025.5762).

Copyright: © 2025 Mujeebulla et al. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, that permits unrestricted use, distribution and reproduction in any medium after the author(s) and source are credited.

Data Availability Statement: Legal restrictions are imposed on the public sharing of raw data. However, authors have full right to transfer or share the data in raw form upon request subject to either meeting the conditions of the original consents and the original research study. Further, access of data needs to meet whether the user complies with the ethical and legal obligations as data controllers to allow for secondary use of the data outside of the original study.

Conflict of interests: The authors have declared that no conflict of interest exists.

RECEIVED on 06th September 2024

RECEIVED in revised form on 15th February 2025

ACCEPTED in final form on 06th March 2025

PUBLISHED on 20th March 2025

1. INTRODUCTION

The poultry business has experienced unprecedented expansion over the last three decades and important components of the agricultural economy. India ranks fourth in total production of poultry meat in the world (Anonymous, 2019). This has occurred because of increased intake of eggs and meat due to their availability and low cost. Poultry outperform other domesticated animals in terms of maintaining a high feed conversion rate is essential for birds to grow more quickly and attain market weight earlier (Diaz et al., 2019). In India, commercial quail farming has the potential to generate a substantial income and job possibilities. Quails are small-sized birds whose rearing system is easy, simple and adapt to almost all types of climates. Indian climate is suitable for raising quail commercially. At present quail has become the third largest avian species in number only next to chickens and ducks in the country. Quail has a carcass made up of 76% of meat, 14% of skin, and 10% of bone. Quail has the highest meat-to-bone ratio among the poultry species (Gecgel et al., 2015).

It has been realized that the natural feed sources like herbs and its extracts overcome the adverse effect of stress on broiler performance (Aderanti et al., 2024 and Yigit et al., 2014). According to (Goodarzi et al., 2013), organosulfur compounds in onions may have boosted broiler performance by increasing nutrient absorption. Onion extract exhibited non-toxic and non-pathogenic effects on animals and considered as feed additive. Antibody production was higher in chicken blood serum if we supplemented onion in their feed. Onion (*Allium cepa*) belongs to the family Amaryllidaceae. It comprises the active principle quercetin, allylic, and allicin sulfide complex, which possesses anticancer, antioxidant, antimicrobial, anti-diabetic, hypocholesterolemic, hypolipidaemic (Muhammad et al., 2023). Factors affecting the impact of natural antioxidants on broiler meat are inclusion level, environmental conditions, basal diet and potential associations with the condition of rearing (Kirkpinar et al., 2014). Onion meal inclusion in chicken diets has been shown to improve gut morphology and carcass quality of poultry meat (Ur Rahman et al., 2017; br Nainggolan et al., 2024, Al-Ramamneh, 2018). Carcass characteristics of meat are the major factors that attract consumers to buy meat (Yang et al., 2005; Tella et al., 2023). In experimental mice with hypercholesterolemia, dramatically lower serum cholesterol levels (Vidyavati et al., 2010). Likewise, rats blood cholesterol levels decreased when supplemented with onions (Baksh et al., 2011). Onions have a hypocholesterolemic impact because they either boost the liver absorption of high-density lipoprotein (HDL) or decrease the production of cholesterol (Goodarzi et al., 2013).

High feed costs which are partially attributable to the price fluctuations of feed ingredients present a major challenge to successful poultry production. This often results in the use of feed ingredients and feed additives that are cheap and of low quality in an effort to improve nutrient exploitation by the bird (Aji et al., 2011; Mammo, 2012; Mulugeta et al., 2019). Adding feed additives to poultry diets maximizes nutrient exploitation by the bird thereby improving growth performance (Frankic et al., 2009) and feed conversion efficiency (Mulugeta et al., 2019).

Onion is most frequently used as powder or extracts (Liguori et al., 2017; Pareek et al., 2017). Onion waste comprising of skin/peel, two external fleshy leaves, top and bottom part, is generated in considerable amounts during industrial processing, and these wastes are rich in bioactive compounds/phytochemicals (Benítez et al., 2011). Keeping in view of the significant importance of OPP and quail meat gaining importance, the current study was proposed to evaluate the effect of supplementation of onion peel powder at varying levels on the carcass traits, meat quality and cost economics of production of Japanese quail.

2. MATERIALS AND METHODS

2.1. Experimental site

The present experiment was planned during November to December, 2023 to study the effect of dietary supplementation of Onion peel powder on the performance of Japanese quail. The biological trial was carried out at the Poultry unit, Livestock Farm Complex, N.T.R. College of Veterinary Science, Gannavaram (A.P). Laboratory analysis was carried out at the Department of Animal Nutrition, N.T.R. College of Veterinary Science, Gannavaram (A.P).

2.2. Procurement of feed ingredients

Onion peels procured from the local market were dried and powdered in Willey's grinder for incorporation into quail diets. Maize, Soybean meal, trace minerals, stone grit, methionine, vitamins, salt, liver tonic, toxin binder, coccidiostat, DCP, and palm oil were procured from the local market and were ground, mixed in the feed mixing plant attached to Department of Animal Nutrition.

2.3. Proximate analysis

The chemical composition of feed ingredients utilized in this experiment and the experimental diets were determined as per Anonymous (2007), while the calcium and phosphorus contents were estimated. The gross energy of Onion Peel powder was estimated using a Bomb Calorimeter and the Metabolizable energy value of Onion Peel powder was derived by using formula. $(ME \text{ (kcal kg}^{-1}) = \text{Gross energy (cal g}^{-1}) \times 0.82)$

2.4. Preparation of basal diets

Experimental diets were prepared as T₁ (Basal diet), T₂ (Basal diet+10 g kg⁻¹ Onion peel powder), T₃ (Basal diet+20 g kg⁻¹ Onion peel powder), T₄ (Basal diet+30 g kg⁻¹ Onion peel powder) and T₅ (Basal diet+40 g kg⁻¹ Onion peel powder). The basal diets were prepared as per the nutrient requirements of poultry (Anonymous, 1994) for Japanese quail.

2.5. Experimental birds and their management

Two hundred-and twenty-five-day-old Japanese quail chicks were weighed individually and randomly divided into five equal groups of three replicates each with 15 chicks per replicate. Each group was allotted to one of the dietary treatments at random. The experiment was conducted from 0–5 weeks of age. All the quail chicks were housed in 5-tier battery cages throughout the experiment. Feed and water were provided *ad libitum*. Feed was offered regularly and the feed left was weighed once a week to quantify the feed intake. B-complex vitamins were added to the water for three days during the first week. The experiment was carried out under uniform management practices for a period of five weeks.

2.6. Data collection

2.6.1. Carcass characteristics

At the end of the trial period (5th week), two birds per replicate and thus a total of six birds per treatment were randomly selected, weighed, and slaughtered. The data on carcass yield, dressing percentage, ready-to-cook yield, and weight of heart, liver, and gizzard were collected, weighed, and recorded. Thus, relative weights (% of live body weight at slaughter) of carcass yield plus total edible organs were calculated.

2.7. Estimation of meat quality parameter

2.7.1. pH

The pH of the meat sample was determined by following the method of Trout et al. (1992) using a deluxe digital pH meter (model 101E). A representative sample of 5 g was homogenized with 45 ml of distilled water in a laboratory blender for about one minute. The pH was recorded by immersing the combination glass electrode of a digital pH meter in the homogenate. Before the measurement of pH, the pH meter was calibrated with buffer solutions of pH 4 and pH 7 as per user manual instructions to avoid errors.

2.8. Extract release volume (ERV)

Homogenize 20 g of meat with 100 ml of distilled water for 2 minutes. Poured the homogenate directly into the funnel lined with Whatman filter paper No.1, folded thrice to make eight sections. Allowed the homogenate to seep between the folds and then collected the extract in a 100 ml graduated

cylinder for 15 minutes.

2.9. Water holding capacity (WHC)

The water-holding capacity of the meat samples was measured according to the method as described by Wardlaw et al. (1973). A 20 g finely chopped meat sample was placed in a centrifuge tube containing 30 ml NaCl (0.6 M) and was stirred with a glass rod for 1 minute. The tube was then kept at refrigeration temperature (4±1°C) for 15 minutes, stirred again, and centrifuged at 3000 rpm using a refrigerated centrifuge for 15 min. The supernatant was measured and the amount of water retained by samples was expressed as WHC in percentage.

Water holding capacity (%)=(Area of expressed water/Area of pressed meat sample)×100

2.10. Sensory evaluation of meat

The meat samples were cooked and subjected to a nine-member taste panel for sensory evaluation of color, appearance, flavor, juiciness, tenderness, and overall acceptability on a nine-point Hedonic scale according to Cross et al. (1978).

2.11. Cost economics

The relative economics of rearing quails up to 5 weeks of age by supplementing Onion peel powder as growth promoter varying levels in the diets was calculated based on the actual ingredient cost at the prevailing market.

2.12. Statistical analysis

Statistical analysis of the data was carried out according to the procedures suggested by Snedecor et al. (1994). The data obtained were subjected to one-way ANOVA. Differences between means were tested at the 5% probability level using Duncans (1955) LSD test.

3. RESULTS AND DISCUSSION

3.1. Pre-slaughter live weight (g)

The mean results of (±S.E) pre-slaughter live weights of Japanese quail reared from day-old to five weeks of age by feeding OPP is significantly higher in the T₅ group compared to all other treatments are presented in table 1.

3.2. Liver weight (%)

The present study revealed that supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ diet from day old to five weeks had no significant effect ($p>0.05$) in liver weights presented in Table 1. In line with the present findings, Goodarzi and Nanekarani (2014), An et al. (2015), Keohavong and Bounyavong (2017), Olusola et al. (2018), Farahani et al. (2015), Aditya et al. (2017) and Omar et al. (2020) reported that Onion extract supplementation had no significant effect ($p>0.05$) on liver weight.

Table 1: Mean ($\pm$ S.E) carcass traits of J. quail fed with varying levels of OPP from day old to five weeks

Treatment (OPP g kg ⁻¹)	Carcass traits						
	Pre slaughter live weight(g)	Liver weight (%)	Heart weight (%)	Gizzard weight (%)	Giblet weight (%)	Carcass weight (g)	Dressing percentage
T ₁ (0)	195.33 ^b $\pm$ 1.73	1.14 ^b $\pm$ 0.05	2.46 $\pm$ 0.09	2.57 $\pm$ 0.27	6.43 $\pm$ 0.22	126.53 ^b $\pm$ 1.94	64.77 $\pm$ 1.13
T ₂ (10)	197.50 ^b $\pm$ 2.99	1.22 ^b $\pm$ 0.07	2.68 $\pm$ 0.23	2.70 $\pm$ 0.15	6.61 $\pm$ 0.32	129.28 ^b $\pm$ 1.31	65.46 $\pm$ 0.29
T ₃ (20)	198.00 ^b $\pm$ 1.76	1.21 ^b $\pm$ 0.05	2.86 $\pm$ 0.14	2.76 $\pm$ 0.12	6.95 $\pm$ 0.29	130.57 ^b $\pm$ 0.77	65.95 $\pm$ 0.30
T ₄ (30)	205.50 ^a $\pm$ 1.23	1.30 ^{ab} $\pm$ 0.06	2.86 $\pm$ 0.35	2.70 $\pm$ 0.20	7.00 $\pm$ 0.47	137.28 ^a $\pm$ 1.75	66.80 $\pm$ 0.99
T ₅ (40)	205.83 ^a $\pm$ 2.58	1.45 ^a $\pm$ 0.06	3.13 $\pm$ 0.24	2.94 $\pm$ 0.08	7.02 $\pm$ 0.19	138.96 ^a $\pm$ 2.62	67.51 $\pm$ 0.73
SEm $\pm$	1.402	0.032	0.104	0.078	0.219	0.858	0.507
N	6	6	6	6	6	6	6
P	0.001	0.020	0.363	0.708	0.607	0.001	0.249
SS	**	*	NS	NS	NS	**	NS

Values in column bearing different super scripts differ significantly ** ($p < 0.01$), * ($p < 0.05$), NS ($p > 0.05$)

Contrary to present findings Goliomytis et al. (2014) reported that 1 g kg⁻¹ Quercetin supplementation resulted in significantly ($p < 0.05$) improved liver weights. Al-Ramamneh et al. (2018) reported that the addition of 2.5% onion to the diet resulted in a substantial ($p < 0.05$) increase in liver weight.

3.3. Heart weight (%)

The present study revealed that supplementation of OPP at (0, 10, 20, 30, and 40) g kg⁻¹ diet from day old to five weeks presented in Table 1, there was no significant difference ($p > 0.05$) in heart weights. Farahani et al. (2015), Aditya et al. (2017), Tella et al. (2023) and Omar et al. (2020) reported that supplementation of Onion extract had no significant ($p > 0.05$) change in heart weight. Contrary to the present findings, Olusola et al. (2018) reported that the Onion skin meal (100 g kg⁻¹) fed group had substantially ($p < 0.05$) increased heart weight than the other treatments. Keohavong and Bounyavong (2017) reported that the supplemented Onion extract group had significantly ($p < 0.05$) lowered heart weight.

3.4. Gizzard weight (%)

The present study revealed that supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ from day old to five weeks had no significant effect ($p > 0.05$) in gizzard weights though numerically higher gizzard weights with an increase in levels of onion peel powder were observed are presented in table 1.

Contrary to the present findings, Keohavong and Bounyavong (2017) and Omar et al. (2020) reported that supplementation of 2.5%, 1%, and 1 g kg⁻¹ Onion extract significantly ($p \leq 0.001$) increased gizzard weight.

3.5. Giblet weight (%)

The giblet weight ranged from 6.43 to 7.02 among different

treatments on supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ from day old to five weeks presented in table 1.

3.6. Carcass weight (g)

The carcass weight ranged from 126.53 to 138.96 g among different treatments on supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ from day old to five weeks are presented in table 1.

3.7. Dressing percentage (%)

The present study revealed that supplementation of varying levels of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ from day old to five weeks presented in table 1, had no significant effect ($p > 0.05$) on dressing percentage. In accordance with the present findings, Farahani et al. (2015) and Omar et al. (2020) reported that there was no significant ($p > 0.05$) difference in dressing percentage by supplementing onion extract. Kazeem et al. (2021) opined that Onion leaf powder supplementation had no significant ($p > 0.05$) difference in dressing percentage. Contrary to the present findings, Keohavong and Bounyavong (2017) reported that supplementing with 1% onion extract was significantly ($p < 0.05$) higher dressing percentage.

There are theories that say essential oils can help with digestion. The fact that herbs and spices (Onion) help in the digestion of feed which increases nutrient absorption and in turn increases body weight and the size of organs can be increased is well established according to previous studies. These might be the reasons for improved carcass traits (Goodarzi and Nanekarani, 2014) on supplementation of OPP.

3.8. pH

The present study revealed that supplementation of OPP in the diet had no effect on the pH value of meat in quail.

Table 2: Mean ($\pm$ S.E) meat pH, ERV, and WHC of Japanese quail fed with varying levels of OPP from day old to five weeks

Treatment (OPP g kg ⁻¹)	pH	ERV	WHC
T ₁ (0)	5.30 $\pm$ 0.38	19 $\pm$ 0.40	23.3 $\pm$ 0.34
T ₂ (10)	5.42 $\pm$ 0.53	21 $\pm$ 0.40	24.3 $\pm$ 0.14
T ₃ (20)	5.30 $\pm$ 0.30	20 $\pm$ 0.30	25.0 $\pm$ 0.11
T ₄ (30)	5.62 $\pm$ 0.31	23 $\pm$ 0.30	22.4 $\pm$ 0.21
T ₅ (40)	5.97 $\pm$ 0.29	25 $\pm$ 0.20	21.4 $\pm$ 0.14
SEm $\pm$	0.330	0.316	0.272
N	6	6	6
P	0.192	0.116	0.325
SS	NS	NS	NS

The pH ranged from 5.30 to 5.97 which is within the normal range for quail meat are presented in table 2. The results were like the findings of Goliomytis et al. (2014) who reported that supplementation of quercetin had no significant effect ($p>0.05$) on pH. An et al. (2015) indicated that there was no significant difference ($p>0.05$) in pH on supplementation of Onion extract, Olusola et al. (2018) reported that supplementation of Onion skin extract and Onion skin meal had no significant ($p>0.05$) difference in pH. Kazeem et al. (2021) reported that supplemented onion leaf powder did not have a significant ($p>0.05$) difference in pH.

The addition of antioxidant supplements has no impact on the pH of broiler meat. This may be because pre-slaughter stress on the animal and post-slaughter handling of the meat affects the pH of the meat, and the pH of the meat may

have indicated varied glycogen reserves before slaughter, according to Simitzis et al. (2008).

3.9. Extract release volume (erv)

ERV ranged from 19 to 25 ml among different treatments on supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ diet from day old to five weeks are presented in table 2.

3.10. Water holding capacity (whc)

WHC content ranged from 21.4 to 25.0% among different treatments on supplementation of OPP at 0, 10, 20, 30, and 40 g kg⁻¹ diet from day old to five weeks are presented in table 2.

3.11. Effect of dietary supplementation of onion peel powder on sensory evaluation of meat

The present study revealed that supplementation of OPP in the diet had no effect ($p>0.05$) on sensory parameters of quail meat but resulted in better color, tenderness, juiciness, and flavor values compared to the control which used to overall acceptability of the quail meat are presented in table 3. According to (br Nainggolan et al., 2024) does not affect the color.

3.12. Cost economics

The feed cost kg⁻¹ gain in quail decreased in all treatment groups as compared to the control. The lowest feed cost kg⁻¹ gain was recorded at 40 g kg⁻¹ OPP are presented in table 4. The decreased feed cost kg⁻¹ gain in the 40 g kg⁻¹ OPP group of quail observed in the present study might be attributed to better feed efficiency and increased weight gains as compared to the other groups. Due to the presence of sulfur compounds and high palatability, increased weight gains on supplementation which in turn leads to high profits.

Similarly, Aji et al. (2011) reported that supplementing 25

Table 3: Mean ($\pm$ S.E) scores for sensory evaluation of meat of Japanese quail fed with varying levels of OPP from day old to five weeks

Treatment (OPP g kg ⁻¹)	Color	Flavor	Juiciness	Tenderness	Overall acceptability
T ₁ (0)	7.33 $\pm$ 0.33	7.17 $\pm$ 0.54	7.33 $\pm$ 0.42	7.67 $\pm$ 0.55	7.17 $\pm$ 0.30
T ₂ (10)	8.00 $\pm$ 0.36	7.50 $\pm$ 0.50	7.50 $\pm$ 0.67	7.33 $\pm$ 0.55	7.50 $\pm$ 0.50
T ₃ (20)	8.17 $\pm$ 0.30	7.83 $\pm$ 0.74	7.67 $\pm$ 0.76	8.00 $\pm$ 0.68	7.83 $\pm$ 0.30
T ₄ (30)	7.83 $\pm$ 0.30	7.67 $\pm$ 0.55	7.87 $\pm$ 0.55	7.50 $\pm$ 0.34	8.00 $\pm$ 0.36
T ₅ (40)	8.33 $\pm$ 0.21	8.17 $\pm$ 0.47	7.83 $\pm$ 0.30	8.83 $\pm$ 0.40	8.33 $\pm$ 0.33
SEm $\pm$	0.143	0.246	0.238	0.238	0.171
N	6	6	6	6	6
P	0.221	0.791	0.977	0.300	0.237
SS	NS	NS	NS	NS	NS

NS-non significant ($p>0.05$)

Table 4: Mean ($\pm$ S.E) cost economics of Japanese quail fed with OPP from day old to five weeks

Treatment (OPP g kg ⁻¹)	Cost economics				
	Cost of feed 100 kg ⁻¹ (₹)	Cumulative feed intake (g)	Cost of feed bird ⁻¹ (₹)	Body weight gain (g)	Feed cost kg ⁻¹ live weight gain (₹)
T ₁ (0)	47.70	630.74	30.08	182.67	164.04
T ₂ (10)	47.80	628.36	30.03	189.62	158.02
T ₃ (20)	47.90	625.35	29.95	189.07	158.10
T ₄ (30)	48.00	624.40	29.97	195.53	153.33
T ₅ (40)	48.10	623.72	30.00	198.00	151.26

1US\$=INR 83.24 (average value of December, 2023)

and 50 mg of onion considerably ($p < 0.05$) reduced feed cost compared to control.

4. CONCLUSION

Increasing OPP levels in Japanese quails significantly increased carcass traits like live weight, heart weight, and carcass weight. However, supplementation of OPP from 0 to 40 g kg⁻¹ in diet did not affect dressing percentage, liver, gizzard, and giblet weights. Meat quality parameters like pH, ERV, and WHC remained unaffected. Sensory evaluations showed no significant differences in color, flavor, juiciness, tenderness, or overall acceptability. Feed cost kg⁻¹ gain was significantly lower in quails fed with 40 g kg⁻¹ OPP.

5. REFERENCES

- Aderanti, I.O., Oluwagbemiga, O.A., 2024. The role of phytochemical feed additives in stress mitigation in broiler chickens. *Journal of Animal Physiology and Animal Nutrition* 108(1), 81–98.
- Aditya, S., Ahammed, M., Jang, S.H., Ohh, S.J., 2017. Effects of dietary onion (*Allium cepa*) extract supplementation on performance, apparent total tract retention of nutrients, blood profile, and meat quality of broiler chicks. *Asian-Australasian Journal of Animal Sciences* 30(2), 229–235.
- Aji, S.B., Ignatius, K., Ado, A.Y., Nuhu, J.B., Abdulkarim, A., Aliyu, U., Gambo, M.B., Ibrahim, M.A., Abubakar, H., Bukar, M.M., Imam, H.M., Numan, P.T., 2011. Effect of feeding onion (*Allium cepa*) and garlic (*Allium sativum*) on some performance characteristics of broiler chickens. *Research Journal of Poultry Science* 4(2), 22–27.
- Al-Ramamneh, D., 2018. Reduce heat stress in broiler by adding onion. *Russian Agricultural Sciences* 44(1), 92–96.
- An, B.K., Kim, J.Y., Oh, S.T., Kang, C.W., Cho, S., Kim, S.K., 2015. Effects of onion extracts on growth performance, carcass characteristics and blood profiles of white mini broilers. *Asian-Australasian Journal of Animal Sciences* 28(2), 247–251.
- Anonymous, 2019. Basic Animal Husbandry Statistics (BAHS), Statistical Division, Department of Animal Husbandry, Dairying and Fisheries, Government of India. Available at <https://dahd.gov.in/schemes/programmes/animal-husbandry-statistics> and Access date: 2024-11-26.
- Anonymous, 2007. Official methods of analysis. 18th edition. Association of Analytical Chemists, Washington. Available at: https://www.researchgate.net/publication/292783651_AOAC_2007 and Access date: 2024-11-26.
- Anonymous, 1994. National Research Council, & Subcommittee on Poultry Nutrition. (1994). Nutrient requirements of poultry. National Academies Press. Available at: <https://nap.nationalacademies.org/catalog/2114/nutrient-requirements-of-poultry-ninth-revised-edition-1994> and Access date: 2024-11-26.
- Baksh, R., Chughtai, M.I.D., 2011. Comparative study of onion and garlic on serum-cholesterol, liver cholesterol, prothrombin time and fecal sterols excretion in male albino-rats. *Journal of the Chemical Society of Pakistan* 7(4), 285.
- Benitez, V., Molla, E., Martin-Cabrejas, M.A., Aguilera, Y., Lopez-Andreu, F.J., Cools, K., Terry, L.A., Esteban, R.M., 2011. Characterization of industrial onion wastes (*Allium cepa* L.): dietary fiber and bioactive compounds. *Plant Foods for Human Nutrition* 66(1), 48–57.
- br Nainggolan, S., Trisna, A., Siregar, G.A.W., 2024. The effect of adding batak onion juice (*Allium chinense* G. Don) on the physical and organoleptic quality of chicken nuggets. *Jurnal Peternakan Integratif* 12(3), 132–39.
- Cross, H.R., Moen, R., Stanfield, M., 1978. Guide lines for training and testing judges for sensory analysis of meat quality. *Food Technology* 32, 48–53.
- Diaz Carrasco, J.M., Casanova, N.A., Fernandez Miyakawa, M.E., 2019. Microbiota, gut health, and chicken productivity; what is the connection? *Microorganisms* 7(10), 374.

- Duncan, D.B., 1955. Multiple range and F-tests. *Biometrics* 11, 1–42.
- Farahani, M., Goodarzi, M., Nanekarani, S., 2015. The effects of aqueous extract of onion on performance and some blood biochemical parameters of the Cobb and Ross broilers. *International Journal of Advanced Biological and Biomedical Research* 3(4), 370–377.
- Frankic, T., Voljc, M., Salobir, J., Rezar, V., 2009. Use of herbs and spices and their extracts in animal nutrition. *Acta Agriculturae Slovenica* 94, 95–102.
- Gegel, U., Yilmaz, I., Gurcan, E.K., Karasu, S., Dulger, G.C., 2015. Comparison of fatty acid composition between female and male Japanese quail meats. *Journal of Chemistry* 2015(1), 569746.
- Goliomytis, M., Tsourekis, D., Simitzis, P.E., Charismiadou, M.A., Hager-Theodorides, A.L., Deligeorgis, S.G., 2014. The effects of quercetin dietary supplementation on broiler growth performance, meat quality, and oxidative stability. *Poultry Science* 93(8), 1957–1962.
- Goodarzi, M., Landy, N., Nanekarani, S., 2013. Effect of onion (*Allium cepa* L.) as an antibiotic growth promoter substitution on performance, immune responses and serum biochemical parameters in broiler chicks. *Health* 5, 1210–1215.
- Goodarzi, M., Nanekarani, S., 2014. Effect of onion extract in drink water on performance and carcass traits in broiler chickens. *IERI Procedia* 8, 107–112.
- Kazeem, D.A., Blessing, M.O., Elizabeth, T.A., Anne, E.E., Tobeckukwu, N.A., 2021. Onion leaf and synthetic additives in broiler diet: impact on splenic cytokines, serum immunoglobulins, cecal bacterial population, and muscle antioxidant status. *Journal of the Science of Food and Agriculture* 101(12), 5245–5255.
- Keohavong, B., Bounyavong, S., 2017. Effects of onion (*Allium cepa* L.) extract on growth performance and carcass characteristics of Muscovy duck. *Souphanouvong University Journal Multidisciplinary Research and Development* 3(1), 229–239.
- Kirkpinar, F., Unlu, H.B., Serdaroglu, M., Turp, G.Y., 2014. Effects of dietary oregano and garlic essential oils on carcass characteristics, meat composition, colour, pH and sensory quality of broiler meat. *British Poultry Science* 55(2), 157–166.
- Liguori, L., Califano, R., Albanese, D., Raimo, F., Crescitelli, A., Di Matteo, M., 2017. Chemical composition and antioxidant properties of five white onions (*Allium cepa* L.) landraces. *Journal of Food Quality*, Article ID: 6873651.
- Mammo, M., 2012. The issue of feed-food competition and chicken production for the demands of foods of animal origin. *Asian Journal of Poultry Science* 6(3), 31–43.
- Muhammad, A., Muhammad, H., Khalid, M., 2023. An insight into anticancer, antioxidant, antimicrobial, antidiabetic and anti-inflammatory effects of quercetin. a review. *Polymer Bulletin* 80, 241–262.
- Mulugeta, M., Worku, Z., Seid, A., Debela, L., 2019. Effect of garlic powder (*Allium sativum*) on performance of broiler chicken. *Livestock Research for Rural Development* 31, 4.
- Omar, A.E., Al-Khalaifah, H.S., Mohamed, W.A.M., Gharib, H.A.S., Osman, A., Al-Gabri, N.A., Amer, S.A., 2020. Effects of phenolic-rich onion (*Allium cepa* L.) extract on the growth performance, behavior, intestinal histology, amino acid digestibility, antioxidant activity, and the immune status of broiler chickens. *Frontiers in Veterinary Science* 7, 582–612.
- Tella, A.K., Olubunmi, O.O., Bello, O.M., 2023. Performance and sensory properties of broilers fed finisher diet supplemented with onion skin extract. *Nigerian Journal of Animal Production* 50(5), 107–112.
- Olusola, O., Kehinde, T., Akanbi Akeem, O., 2018. Performance and meat quality attributes of broiler chickens fed onion skin extract and onion skin meal supplementation diets at the finisher stage. *Journal of Experimental Agriculture International* 24(1), 1–7.
- Pareek, S., Sagar, N.A., Sharma, S., Kumar, V., 2017. Onion (*Allium cepa* L.). fruit and vegetable phytochemicals: chemistry and human health. *Hoboken*, 1145–1162.
- Simitzis, P.E., Deligeorgis, S.G., Bizelis, J.A., Dardamani, A., Theodosiou, I., Fegeros, K., 2008. Effect of dietary oregano oil supplementation on lamb meat characteristics. *Meat Science* 79(2), 217–223.
- Snedecor, G.W., Cochran, G.W., 1994. *Statistical methods*, 8th Edition, Iowa State University Press, Ames.
- Trout, E.S., Hunt, N.C., Jhonson, D.E., Claus, J.R., Kastner, C.L., Kropf, D.H., 1992. Chemical, physical and sensory characterization of ground beef containing 5 to 30% fat. *Journal of Food Science* 57, 25–29.
- Ur Rahman, S., Khan, S., Chand, N., Sadique, U., Khan, R.U., 2017. *In vivo* effects of *Allium cepa* L. on the selected gut microflora and intestinal histomorphology in broiler. *Acta Histochemical* 119(5), 446–450.
- Viduyavati, H.G., Manjunatha, H., Hemavathy, J., Srinivasan, K., 2010. Hypolipidemic and antioxidant efficacy of dehydrated onion in experimental rats. *Journal of food Science and Technology* 47, 55–60.
- Wardlaw, F.B., McCaskill, L.H., Acton, J.C., 1973. Effect of postmortem muscle changes on poultry meat loaf properties. *Journal of Food Science* 38(3), 421–423.
- Yang, N., Jiang, R.S., 2005. Recent advances in breeding for quality chickens. *World's Poultry Science Journal* 61(3), 373–381.
- Yigit, A.A., Panda, A.K., Cherian, G., 2014. The avian embryo and its antioxidant defence system. *World's Poultry Science Journal* 70(3), 563–574.