




Effect of Phytobiotic Feed Additives Garlic (*Allium sativum*), Ashwagandha (*Withania somnifera*) and Shatavari (*Asparagus racemosus*) as Alternatives to Antibiotic on Feed Intake, Metabolizability of Nutrients and Balance Study in Broiler Chicks

Hemant Kumar Jediya¹, Monika Joshi²  and S. K. Sharma³

¹Dept. of Animal Nutrition, ²Dept. of Animal Nutrition, ³Dept. of Veterinary Medicine, College of Veterinary and Animal Science, Navania, Udaipur, Rajasthan University of Veterinary and Animal Sciences, Bikaner, Rajasthan (313 601), India



Corresponding  drmonikashivsharma@gmail.com

 0009-0007-2801-0891

ABSTRACT

An experiment was conducted during March to April, 2021 at Poultry unit of College of Veterinary and Animal Science, Navania, Vallabhnagar, Udaipur (Rajasthan), India planned to investigate the effect of phytobiotic feed additives viz., garlic, ashwagandha and shatavari as alternatives to antibiotic on feed intake, metabolizability of nutrients and balance study of broiler chicks. A total of 300 day-old broiler chicks (Cobb-400) were involved in the 42-days. The T₁ i.e. control group was fed on basal diet, while T₂ was supplemented with Oxy tetra cycline (OTC) powder @ 0.1 g kg⁻¹ feed. T₃ and T₄ were supplemented with Garlic powder @ 0.75% and @ 1.50%. T₅ and T₆ were supplemented with Ashwagandha powder @ 0.75% and @ 1.50%. T₇ and T₈ were supplemented with Shatavari powder @ 0.75% and @ 1.50%. T₉ was supplemented with Garlic powder @ 0.25%, Ashwagandha powder @ 0.25% and Shatavari powder @ 0.25%. T₁₀ was supplemented with Garlic powder @ 0.50%, Ashwagandha powder @ 0.50% and Shatavari powder @ 0.50%. Feed intake was a highly significant ($p < 0.01$) effect in different treatments groups. Non significant ($p > 0.05$) effects were observed on dry matter and Organic matter metabolizability in different treatment groups. The metabolizability of crude protein, crude fibre and total ash was highly significant ($p < 0.01$) effect and ether extract had a significant ($p < 0.05$) effect due to treatments groups. The balance study of nitrogen, calcium and phosphorus had a highly significant ($p < 0.01$) effect in different treatment groups.

KEYWORDS: Broiler, feed, garlic, ashwagandha, shatavari powder, metabolizability, nutrients balance

Citation (VANCOUVER): Jediya et al., Effect of Phytobiotic Feed Additives Garlic (*Allium sativum*), Ashwagandha (*Withania somnifera*) and Shatavari (*Asparagus racemosus*) as Alternatives to Antibiotic on Feed Intake, Metabolizability of Nutrients and Balance Study in Broiler Chicks. *International Journal of Bio-resource and Stress Management*, 2025; 16(3), 01-07. [HTTPS://DOI.ORG/10.23910/1.2025.5813](https://doi.org/10.23910/1.2025.5813).

Copyright: © 2025 Jediya et al. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, that permits unrestricted use, distribution and reproduction in any medium after the author(s) and source are credited.

Data Availability Statement: Legal restrictions are imposed on the public sharing of raw data. However, authors have full right to transfer or share the data in raw form upon request subject to either meeting the conditions of the original consents and the original research study. Further, access of data needs to meet whether the user complies with the ethical and legal obligations as data controllers to allow for secondary use of the data outside of the original study.

Conflict of interests: The authors have declared that no conflict of interest exists.

RECEIVED on 25th September 2024

RECEIVED in revised form on 19th February 2025

ACCEPTED in final form on 28th February 2025

PUBLISHED on 05th March 2025

1. INTRODUCTION

Antibiotics have been used widely to prevent infections and poultry diseases and for the improvement of meat and egg production. However, the use of antibiotics is restricted due to drug resistance in bacteria, drug residue in carcasses and also alteration of natural gut micro flora. Kabir et al. (2004) and Hind et al. (2014) found significantly higher levels of diverse antibiotic residues in the edible tissues of chickens given therapeutic or prophylactic antibiotics. For these reasons, many countries around the world have either limited or prohibited the use of antibiotics as growth promoters in livestock. This has created a gap in which animal nutritionists are working intensively to explore safe alternatives from natural sources (Abdel-Aziz et al., 2015; Abu Hafsa et al., 2016). Phytogenic substances are generally regarded as safe in food and feed industries (Elghalid et al., 2020; Kholif and Olafadehan, 2021). Phytogenic compounds have potential antimicrobial effects with multiple mechanisms of action (Dalle Zotte et al., 2016). Thus, traditional herbal knowledge heritage includes uses of turmeric, ginger, garlic, Ashwagandh, Shatavari onion, lemongrass Aloe vera, amla etc. in human and animal nutrition. (Patel and Srinivasan, 2004) have suggested that phytogenic substances can stimulate digestive secretions such as saliva and bile. They reported that improving enzyme activity is the main mode of nutritional action of phytogenic feed additives. Garlic (*Allium sativum*) having the active ingredient alicine, diallyl disulfide has been considered a wonder drug in the herbal world and used as a growth promoter. It improves nutrient utilization and feed conversion efficiency, and improves digestion and immunity. Garlic acts as only a mild antibiotic in comparison to modern antibiotics, but the added advantage of its use as an antibiotic is that the microorganisms do not develop a resistance against garlic as they do against conventional antibiotics. Ashwagandha (*Withania somnifera*), which in turn may improve the performance of birds. Ashwagandha root is bitter to taste and contains several alkaloids (0.13 to 4.30%), which offer medicinal usage. It contains many active principles such as withanolides, withanone, somnitalglucose, rutinoides, inorganic salt and di-hydroxy kaempferol-3 (Pal et al., 2012). These active principles have been reported to possess immunomodulatory, general tonic, hepato-protective, anti-stress, growth promoter and antioxidant properties (Ansari et al., 2008; Singh et al., 2010; Kushwaha et al., 2012 and Varma et al., 2012) beside antibacterial and anti-fungal properties (Punetha et al., 2010). Shatavari possesses nutritive, antistress, adaptogenic, immunomodulatory, galactagogue, anabolic and performance-enhancing properties and is used in various medicinal preparations Shatavari is the one of the most commonly used herbs in traditional medicine due to the presence of steroidal saponins and sapogenins in various

part of the plant (Krishana et al., 2005). The root powder of Shatavari (*Asparagus racemosus*) is used as an herbal feed supplement in poultry feed. Shatavari augment the appetite and stimulates the liver function.

Therefore, considering the above facts the present research was planned to assess the effect of phytobiotic feed additives-garlic (*allium sativum*), ashwagandha (*withania somnifera*) and shatavari (*asparagus racemosus*) as alternatives to antibiotic on feed intake, metabolizability of nutrients and balance study in broiler chicks.

2. MATERIALS AND METHODS

2.1. Study sites

The experiment was conducted during March to April, 2021 at the Poultry unit of the College of Veterinary and Animal Science, Navania, Vallabhnagar, Udaipur (Rajasthan), India.

2.2. Experimental animals and feeding

An experiment was conducted for six weeks on three hundred, day old chicks, unsexed, apparently healthy broiler chicks (VENCOBB). Each bird was weighed individually on arrival and randomly distributed to ten different dietary treatments groups ($T_1 - T_{10}$) using a completely randomized design (CRD). The body weight of chicks has remained similar in all the groups. Each random treatment has three replicates ($R_1 - R_3$) having 10 birds in each replicate. The ISO certified basal feed in the form of broiler pre starter, broiler starter and broiler finisher was procured from feed distributor "Udaipur Kukkut Utpadak Sahkari Samiti Ltd.", Udaipur (Rajasthan) in required amount. Phytobiotic feed additives garlic (*allium sativum*), ashwagandha (*withania somnifera*) and shatavari (*asparagus racemosus*) powders were procured from the local market. The Proximate composition of the experimental feed was analyzed according to standard methods of analysis (Anonymous, 2016). The chemical composition of the Experimental feed is presented in (Table 1). The feeding was done in three phases, broiler pre starter (0–7 days), broiler starter (8–21 days) and broiler finisher (22–42 days) feeds for different treatments as per the guidelines of (Anonymous, 2007). Diet T_1 served as control while diet T_2 was supplemented with Oxy tetracycline (OTC) powder @ 0.1 g kg⁻¹ feed. T_3 and T_4 were served as Basal diet supplemented with Garlic powder @ 0.75% and @ 1.50%. T_5 and T_6 were served as Basal diet supplemented with Ashwagandha root powder @ 0.75% and @ 1.50%. T_7 and T_8 were served as Basal diet supplemented with Shatavari root powder @ 0.75% and @ 1.50%. T_9 was served as a Basal diet supplemented with Garlic powder @ 0.25%, Ashwagandha root powder @ 0.25% and Shatavari root powder @ 0.25%. T_{10} was served as a Basal diet supplemented with Garlic powder @ 0.50%, Ashwagandha root powder @ 0.50% and Shatavari root powder @ 0.50%.

Table 1: Proximate composition of broiler pre starter, broiler starter Broiler finisher, *Allium sativum* (garlic) powder, *Withania somnifera* (Ashwagandha) root powder and *Asparagus racemosus* (Shatavari) root powder

| Chemical composition | Broiler pre starter | Broiler starter | Broiler finisher | <i>Allium sativum</i> | <i>Withania somnifera</i> | <i>Asparagus racemosus</i> |
|---------------------------------|---------------------|-----------------|------------------|-----------------------|---------------------------|----------------------------|
| <u>Proximate principles (%)</u> | | | | | | |
| Dry matter | 89.72 | 91.22 | 91.52 | 91.52 | 94.60 | 90.99 |
| <u>Organic</u> | | | | | | |
| Matter | 95.53 | 93.23 | 93.28 | 96.84 | 94.32 | 96.00 |
| Crude protein | 23.09 | 22.38 | 20.29 | 18.20 | 5.21 | 15.29 |
| Ether extract | 2.93 | 4.26 | 4.68 | 5.32 | 0.57 | 0.66 |
| Crude fibre | 4.57 | 3.74 | 3.79 | 3.12 | 12.89 | 2.82 |
| Total ash | 4.47 | 6.77 | 6.72 | 3.16 | 5.68 | 4.00 |
| <u>Nitrogen free</u> | | | | | | |
| Extract | 64.94 | 62.85 | 64.52 | 70.20 | 75.65 | 77.23 |
| <u>Mineral composition (%)</u> | | | | | | |
| Calcium | 1.07 | 1.09 | 1.06 | 0.50 | 1.17 | 0.24 |
| Phosphorus | 0.69 | 0.70 | 0.63 | 0.70 | 0.63 | 0.86 |

2.3. Method of data collection

A metabolism trial of 7 days was conducted at the end of the experiment. The birds were fed individually with their respective treatment diets. The adaptation period was approximately 2 days, followed by a collection period of 5 days. During the collection period, data on the quantity of feed offered, left over and excreta voided were recorded in order to determine the feed intake, nutrients utilization and nitrogen, calcium and phosphorus retention. The daily feed intake was calculated after the deduction of the leftovers from the feed offered. The digestibility/metabolizability of nutrients was analysed by proximate principles as per (Anonymous, 2016). The total nitrogen content of feed and excreta was determined through Kjeldahl's method using Kel plus Automatic Nitrogen Analyzer equipment. The calcium and phosphorus of feed and excreta samples was estimated using procedures described by (Talpatra et al., 1940). The per cent nitrogen, calcium and phosphorus retention under different treatments was calculated through percent difference in intake and outgo of nutrients. The data obtained in the experiment were analyzed using analysis of variance (ANOVA) by (Snedecor and Cochran, 2004). Significance of means differences were tested by Duncan's New Multiple Range Test (DNMART) (Duncan, 1955) as modified by (Kramer, 1957).

3. RESULTS AND DISCUSSION

3.1. Average feed intake

Average weekly feed intake by the broilers in six weeks period in T₁ (Control), T₂, T₃, T₄, T₅, T₆, T₇, T₈, T₉ and

T₁₀ treatment groups were found to be 4352.78, 4178.46, 4119.18, 4082.93, 3946.17, 3917.96, 3936.27, 3937.93, 3966.52 and 3957.94 g, respectively (Table 2). The highest average feed intake was found in T₁ (Control) group whereas, lowest average feed intake was found in T₆ group i.e., broiler fed a basal diet supplemented with ashwagandha root powder @ 1.50%. The statistical analysis of data revealed highly significant ($p < 0.01$) effect in terms of average weekly feed intake with dietary supplementation of garlic, Ashwagandha and Shatavari alone and in combination at different levels in the ration of broiler chicks. The present findings are in agreement with Eid et al. (2014); Fayed et al. (2011); Abdullah et al. (2010); Raeesi et al. (2010); Abou-Elangana et al. (2016); Jimoh et al. (2013); Samanthi et al. (2015) who reported significant decrease in feed consumption due to supplementation of Garlic powder probably due to the associated flavor factor. Shisodiya et al. (2008); Muhammad et al. (2009); Bhardwaj et al. (2011); Abdallah et al. (2016) and Mane et al. (2012); Pandey et al. (2013); Gaikwad et al. (2015); Mali et al. (2017) reported that significant decrease in feed consumption due to supplementation of Ashwagandha and Shatavari powder as compare to the control group.

3.2. Metabolizability of nutrients

The mean values of dry matter metabolizability (%) and organic matter metabolizability (%) were found to be 73.12 and 76.03, 73.74 and 76.81, 74.03 and 77.19, 74.61 and 77.68, 74.87 and 77.83, 75.30 and 78.21, 75.22 and 78.03, 75.43 and 78.42, 76.02 and 78.83, 76.16 and 79.21 in T₁ (Control), T₂, T₃, T₄, T₅, T₆, T₇, T₈, T₉ and T₁₀ treatment

Table 2: Effect of Garlic powder, Ashwagandha root powder and Shatavari root powder on feed intake (g bird⁻¹ week⁻¹) in broiler chicks

| | Treatment groups | | | | | | | | | | SEm |
|----------|----------------------|----------------------|-----------------------|-----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|--------|
| | T ₁ | T ₂ | T ₃ | T ₄ | T ₅ | T ₆ | T ₇ | T ₈ | T ₉ | T ₁₀ | |
| I Week | 162.26 | 161.33 | 159.36 | 159.78 | 161.74 | 161.76 | 160.24 | 158.18 | 164.96 | 165.98 | 3.249 |
| II Week | 367.03 ^c | 364.26 ^c | 416.14 ^a | 413.98 ^a | 399.87 ^a | 396.89 ^{ab} | 403.81 ^a | 398.47 ^a | 402.94 ^a | 403.47 ^a | 7.789 |
| III Week | 670.96 | 700.33 | 714.96 | 708.94 | 690.19 | 694.56 | 694.14 | 695.19 | 699.72 | 690.49 | 12.074 |
| IV Week | 768.32 | 770.06 | 776.12 | 772.12 | 766.36 | 767.34 | 764.98 | 764.88 | 765.76 | 766.44 | 15.483 |
| V Week | 952.14 | 908.34 | 968.46 | 961.92 | 932.14 | 930.87 | 931.37 | 930.35 | 934.47 | 932.38 | 19.964 |
| VI Week | 1432.07 ^a | 1274.14 ^b | 1084.14 ^c | 1066.19 ^c | 995.87 ^d | 966.54 ^d | 981.73 ^d | 990.86 ^d | 998.67 ^d | 999.18 ^d | 20.381 |
| CFI | 4352.78 ^a | 4178.46 ^b | 4119.18 ^{bc} | 4082.93 ^{cd} | 3946.17 ^c | 3917.96 ^c | 3936.27 ^c | 3937.93 ^c | 3966.52 ^c | 3957.94 ^c | 30.304 |

Cumulative feed intake; Means with different superscripts in a row differ significantly

groups respectively and non significant ($p>0.05$) effect were found on dry matter and Organic matter metabolizability. The mean values of crude protein, crude fibre and Total Ash (TA) metabolizability (%) were found to be 74.19, 26.73, and 55.07, 76.22, 27.91 and 55.46, 76.92, 28.47 and 56.39, 77.24, 29.24 and 57.45, 77.84, 30.11 and 58.29, 77.13, 31.07 and 60.21, 78.35, 30.61 and 59.92, 78.63, 31.69 and 61.48, 80.11, 32.78 and 62.71, 83.70, 33.12 and 63.26 in T₁ (Control), T₂, T₃, T₄, T₅, T₆, T₇, T₈, T₉ and T₁₀ treatment groups, respectively (Table 3) and highly significant ($p<0.01$) effect on crude protein, crude fibre and Total Ash (TA) metabolizability. The mean values of ether extract metabolizability (%) was found to be 74.09, 74.88, 75.91, 76.83, 77.04, 78.14, 77.29, 79.33, 80.17 and 81.28 in T₁ (Control), T₂, T₃, T₄, T₅, T₆, T₇, T₈, T₉ and T₁₀ treatment groups, respectively and significant ($p<0.05$) effect on ether extract metabolizability due to supplementation of garlic,

Ashwagandha and Shatavari alone and in combination at different levels in the ration of broiler chicks. Saini, et al. (2017); Kehinde et al. (2019) reported that numerical improvement in dry matter digestibility in the Ashwagandha powder supplemented group as compared to control group. The present study was in agreement with Attia et al. (2017) who reported that the digestibility coefficient of crude protein and ether extract were significantly ($p<0.01$) increased in Ashwagandha powder supplemented group as compared to the control group.

3.3. Balance study

The mean values of nitrogen, calcium and phosphorus balance were found to be 2.28, 0.40 and 0.35, 2.30, 0.43 and 0.37, 2.35, 0.50 and 0.39, 2.37, 0.53 and 0.42, 2.42, 0.55 and 0.41, 2.49, 0.62 and 0.45, 2.45, 0.60 and 0.41, 2.53, 0.63 and 0.47, 2.57, 0.64 and 0.47, 2.60, 0.67 and 0.49 g

Table 3: Effect of garlic powder, ashwagandha root powder and shatavari root powder on metabolizability of nutrients (%) in broiler chicks

| | Treatment groups | | | | | | | | | | SEm |
|----|--------------------|---------------------|---------------------|---------------------|--------------------|---------------------|---------------------|--------------------|---------------------|--------------------|-------|
| | T ₁ | T ₂ | T ₃ | T ₄ | T ₅ | T ₆ | T ₇ | T ₈ | T ₉ | T ₁₀ | |
| DM | 73.12 | 73.74 | 74.03 | 74.61 | 74.87 | 75.30 | 75.22 | 75.43 | 76.02 | 76.16 | 1.144 |
| OM | 76.03 | 76.81 | 77.19 | 77.68 | 77.83 | 78.21 | 78.03 | 78.42 | 78.83 | 79.21 | 1.398 |
| CP | 74.19 ^c | 76.22 ^c | 76.92 ^{bc} | 77.24 ^b | 77.84 ^b | 77.13 ^b | 78.35 ^b | 78.63 ^b | 80.11 ^{ab} | 83.70 ^a | 1.309 |
| EE | 74.09 ^b | 74.88 ^b | 75.91 ^b | 76.83 ^b | 77.04 ^b | 78.14 ^{ab} | 77.29 ^b | 79.33 ^a | 80.17 ^a | 81.28 ^a | 1.235 |
| CF | 26.73 ^d | 27.91 ^{cd} | 28.47 ^c | 29.24 ^{bc} | 30.11 ^b | 31.07 ^{ab} | 30.61 ^b | 31.69 ^a | 32.78 ^a | 33.12 ^a | 0.709 |
| TA | 55.07 ^c | 55.46 ^c | 56.39 ^{bc} | 57.45 ^b | 58.29 ^b | 60.21 ^a | 59.92 ^{ab} | 61.48 ^a | 62.71 ^a | 63.26 ^a | 1.244 |

Means with different superscripts in a row differ significantly

bird⁻¹ day⁻¹ in T₁ (Control), T₂, T₃, T₄, T₅, T₆, T₇, T₈, T₉ and T₁₀ treatment groups, respectively (Table 4). The balance of nitrogen, calcium and phosphorus showed highly significant ($p < 0.01$) effect on with dietary supplementation of garlic, Ashwagandha and Shatavari alone and in combination at different levels as alternatives to Antibiotic growth promoters in the ration of broiler chicks. The present

findings were similar to Saini et al. (2017) reported that significantly increased N balance in broilers fed different levels of Ashwagandha powder as compared to the control group. Ashwagandha contains withanine and withanolides, these compounds may stimulate the gastrointestinal tract secretions which help to increase nutrient digestion and absorption (Jyotsana and Berwal, 2019).

Table 4: Effect of garlic powder, ashwagandha root powder and shatavari root powder on balance of nitrogen, calcium and phosphorus (g bird⁻¹ day⁻¹) in broiler chicks

| | T ₁ | T ₂ | T ₃ | T ₄ | T ₅ | T ₆ | T ₇ | T ₈ | T ₉ | T ₁₀ | SEm |
|--------------------|-------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|-------------------|-------|
| Nitrogen balance | 2.28 ^c | 2.30 ^c | 2.35 ^{de} | 2.37 ^d | 2.42 ^{cd} | 2.49 ^{bc} | 2.45 ^c | 2.53 ^{ab} | 2.57 ^a | 2.60 ^a | 0.024 |
| Calcium balance | 0.40 ⁱ | 0.43 ^h | 0.50 ^g | 0.53 ^{ef} | 0.55 ^e | 0.62 ^{bc} | 0.60 ^{cd} | 0.63 ^b | 0.64 ^b | 0.67 ^a | 0.006 |
| Phosphorus balance | 0.35 ^g | 0.37 ^{fg} | 0.39 ^{ef} | 0.42 ^d | 0.41 ^{de} | 0.45 ^{bc} | 0.41 ^{de} | 0.47 ^{ab} | 0.47 ^{ab} | 0.49 ^a | 0.005 |

Means with different superscripts in a row differ significantly

4. CONCLUSION

The highly significant ($p < 0.01$) effect on average feed intake and balance of nitrogen, calcium and phosphorus. Non significant effect on ($p > 0.05$) dry matter and Organic matter, a significant ($p < 0.05$) on ether extract and highly significant ($p < 0.01$) on crude protein, crude fibre, total ash metabolizability. It is concluded that supplementation of garlic, Ashwagandha and Shatavari alone and in combination at different levels as alternatives to Antibiotic growth promoters showed positive effect on feed intake, Metabolizability and nutrients balance in broiler chicks.

5. REFERENCES

- Abou-Elnaga, M.K., Zanaty, G.A., Abou-Ashour, A.M.H., 2016. Beneficial effects of garlic and thyme powder on performance and carcass characteristics of broiler chicks. *Egyptian Journal of Nutrition and Feeds* 19(2), 363–374.
- Abdel-Aziz, N.A., El-Adawy, M., Mariezcurrena-Berasain, M.A., Salem, A.Z., Olivares-Pérez, J., Kholif, A.E., Borhami, B.E., 2015. Effects of exogenous enzymes, *Lactobacillus acidophilus* or their combination on feed performance response and carcass characteristics of rabbits fed sugarcane bagasse. *Journal of Integrative Agriculture* 14(3), 544–549.
- Abdullah, Y., Mahmoud, K.Z., Nusairat, B.M., Qudsieh, R.I., 2010. Small intestinal histology, production parameters, and meat quality as influenced by dietary supplementation of garlic. *Italian Journal of Animal Science* 9, 80.
- Abdallah, O.A., Omnia, E., Killany, H., El-gharib, E., Raghda, F., Mohamed, 2016. Hematological and Growth Performance Studies after *Withania somnifera* Supplementation in Broilers. *Suez Canal Veterinary Medical Journal* XXI(1), 173–183.
- Abu Hafs, S.H., Salem, A.Z.M., Hassan, A.A., Kholif, A.E., Elghandour, M.M.Y., Barbabosa, A., Lopez, S., 2016. Digestion, growth performance, and caecal fermentation in growing rabbits fed diets containing foliage from browse trees. *World Rabbit Science* 24(4), 283–293.
- Ansari, J.Z., Haq, A., Yousaf, M., Ahmad, T., Khan, S., 2008. Evaluation of different medicinal plants as growth promoters for broiler chicks. *Sarhad Journal of Agriculture* 24(2), 323–330.
- Attia, G., El-Eraky, W., Hassanein, E., El-Gamal, M., Farahat, M., Santana, A.H., 2017. Effect of dietary inclusion of a plant extract blend on broiler growth performance, nutrient digestibility, Cecal microflora and intestinal histomorphology. *International Journal Poultry Sciences* 16, 344–353.
- Anonymous, 2016. Official method of analysis, 20th edition, Association of Official Analytical Chemists, Benjamin Franklin Station, Washington, D.C. Available at: <https://www.scirp.org/reference/referencespapers?referenceid=2502537>.
- Anonymous, 2007. BIS, 2007. Poultry Feeds Specification: IS: 1374 (5th Revision). Bureau of Indian Standards, Manak Bhavan, 9 Bahadur Shah Zafar Marg, New Delhi-110002, India. Available at: <https://www.bis.gov.in/wp-content/uploads/2023/01/Product-ManualPM-2-for-IS-1374.pdf>.
- Bhardwaj, R.K., Gangwar, S.K., 2011. Effect of dietary supplementation of *Withania somnifera* on egg production and egg quality parameters in Japanese quails. *International Journal of Advanced Biological Research* 1(1), 32–34.
- Dalle Zotte, A., Celia, C., Szendro, Z., 2016. Herbs and spices inclusion as feedstuff or additive in growing rabbit diets and as additive in rabbit meat: A

- review. *Livestock Science* 189, 82–90.
- Duncan, D.B., 1955. Multiple range test. *Biometrics* 11, 1–6.
- Eid, K.M., Iraqi, M.M., 2014. Effect of garlic powder on growth performance and immune response for newcastle and avian influenza virus diseases in broiler of chickens. *Animal Biotechnology Poultry and Fish* 1, 7–13.
- Elghalid, O.A., Kholif, A.E., El-Ashry, G.M., Matloup, O.H., Olafadehan, O.A., El-Raffa, A.M., Abd El-Hady, A.M., 2020. Oral supplementation of the diet of growing rabbits with a newly developed mixture of herbal plants and spices enriched with special extracts and essential oils affects their productive performance and immune status. *Livestock Science* 238, 104082. <https://doi.org/10.1016/j.livsci.2020.104082>.
- Fayed, R.H., Abeer, H., Razek, A., Jehan, M., 2011. Effect of dietary garlic supplementation on performance, carcass traits, and meat quality in broiler chickens. *Parameters* 100, 1000–1004.
- Gaikwad, D.S., Nage, S.P., Chavan, S.D., Jadhav, D.B., 2015. Effect of supplementation of Shatavari root powder on broiler characteristics and economics of Vencobb–400 chicks. *The Ecoscan* 7, 259–262.
- Hind, A.E., Adil, M., Samah, A., 2014. Screening of antibiotic residues in poultry liver, kidney and muscle in Khartoum State, Sudan. *Journal of Applied and Industrial Sciences* 2(3), 116–122.
- Jimoh, A.A., Ibitoye, E.B., Dabai, Y.U., Garba, S., 2013. *In vivo* antimicrobial potentials of garlic against *Clostridium perfringens* and its promotant effects on performance of broiler chickens. *Pakistan Journal of Biological Sciences* 16(24), 1978–1984.
- Jyotsana, P.K., Berwal, R., 2019. Supplementation of Ashwagandha root powder improving gut micro flora and immune response of broilers. *The Pharma Innovation Journal* 8(10), 82–85.
- Kabir, J., Umoh, V.J., Audu-Okoh, E., Umoh, J.U., Kwaga, J.K.P., 2004. Veterinary drug use in poultry farms and determination of antimicrobial drug residues in commercial eggs and slaughtered chicken in Kaduna State, Nigeria. *Food Control* 15(2), 99–105.
- Kehinde, A.S., Babatunde, T.O., Kehinde, O.J., 2019. Effect of garlic supplementation in the diets of cockerel chicks on performance and economy of production. *Asian Journal of Agricultural Extension, Economics and Sociology* 35 (4), 1–7.
- Kholif, A.E., Olafadehan, O.A., 2021. Essential oils and phytogenic feed additives in ruminant diet: Chemistry, ruminal microbiota and fermentation, feed utilization and productive performance. *Phytochemistry Reviews* 20(6), 1087–1108.
- Kramer, C.Y., 1957. Extension of multiple range tests to group means with unequal numbers of replications. *Biometrics* 12, 307–310.
- Krishana, L., Swarup, D., Patra, R.C., 2005. An overview prospects of ethano-veterinary medicine in India. *Indian Journal of Animal Science* 75(12), 1481–1491.
- Kushwaha, S., Agatha, B., Chawla, P., 2012. Effect of ashwagandha (*Withania somnifera*) root powder supplementation in treatment of hypertension. *Journal of Ethnobiology and Ethnomedicine* 6, 111–115.
- Mali, R.G., Hegokar, S.R.S., Adangle, S.B., Walkunde, T.R., 2017. Effect of supplementation of ashwagandha (*Withania somnifera*) and shatavari (*Asparagus racemosus*) on growth performance of Giriraja poultry birds. *Trends in Biosciences* 10(36), 7558–7562.
- Mane, A.G., Kulkarni, A.N., Korake, R.L., Bhutkar, S.S., 2012. Effect of supplementation of ashwagandha (*Withania somnifera*) and shatavari (*Asparagus racemosus*) on growth performance of broilers. *Research Journal of Animal Husbandry and Dairy Science* 3(2), 94–96.
- Muhammad, J., Durrani, F., Hafeez, A., Khan, R.U., Ahmad, I., 2009. Effect of aqueous extract of plant mixture on carcass quality of broiler chicks. *Research Journal of Agriculture and Biological Sciences* 4(1), 37–39.
- Pal, A., Mahadeva, N., Farhath, K., Bawa, A.S., 2012. *In-vitro* studies on the antioxidant assay profiling of root of *Withania somnifera* L. (Ashwagandha) Dunal. *Agriculturae Conspectus Scientificus* 77, 95–101.
- Patel, K., Srinivasan, K., 2004. Digestive stimulant action of spices: a myth or reality? *Indian Journal of Medical Research* 119, 167–179.
- Pandey, N.K., Singh, D.P., Niwas, R., 2013. Broiler characteristics, sensory qualities, and economic efficiency in vencobb–400 chicks supplemented with a conjugated herbal feed additive in diet. *Animal Science Reporter* 7(4), 128–132.
- Punetha, A., Muthukumar, J., Hemrom, A.J., Arumugam, N., Jayakanthan, M., Sundar, D., 2010. Towards understanding the regulation of rubber biosynthesis: Insights into the initiator and elongator enzymes. *Journal of Bioinformatics and Sequence Analysis* 2, 1–10.
- Raeesi, M., Hoseini-Aliabad, S.A., Roofchae, A., Zare Shahneh, A., Pirali, S., 2010. Effect of periodically use of garlic (*Allium sativum*) powder on performance and carcass characteristics in broiler chickens. *World Academy of Science, Engineering and Technology* 68, 1213–1219.
- Samanthi, K.A.M., Nayananjali, W.A.D., Adikari, A.M.J.B., Liyanage, R., 2015. Dietary garlic (*Allium sativum* L.) supplementation on performance, meat

- quality and lipid profile in broilers. Rajarata University Journal 3, 17–24.
- Saini, J., Sharma, T., Dhuria, R.K., Kumari, N., Mitharwal, N., Limba, A.K., Meena, M., 2017. Effect of feeding *Withania somnifera* and enzyme alone and in Combination on dry matter digestibility and nitrogen balance of Broiler chicks. Veterinary Practitioner 18(1), 124–125.
- Singh, B., Chandan, B.K., Gupta, D.K., 2010. Chemistry of garlic (*Allium sativum*) with special reference to allin and allicin-A review. In: Effect of dietary inclusion of plant extracts on the growth performance of male broiler. Poultry Science (Suppl), 43–44.
- Shisodiya, J.M., Chpoad, S.S., Rajput, A.B., Chandankhede, J.M., Ingale, K.S., Kolte, B.R., 2008. Comparative study of ashwagandha and commercial synthetic compound on performance of broilers during hot weather. Veterinary World 1(10), 310–311.
- Snedecor, G.W., Cochran, W.G., 2004. Statistical methods. 8th Edn., Oxford and IBH publishing company, Kolkata.
- Talpatra, S.K., Ray, S.C., Sen, K.C., 1940. The analysis of mineral constituents in biological material. Estimation of phosphorus, chloride, calcium, magnesium, sodium and potassium in food stuffs. Journal of Veterinary Science and Animal Husbandry 10, 243–258.
- Varma, N., Kataria, M., Kumar, K., Saini, J., 2012. Comparative study of L asparaginase from different cytotypes of *Withania somnifera* (L.) Dunal and its purification. Indian Journal of Natural Products and Resources 2, 475–481.