



## Effect of Organic Nutrient Sources in Tea [*Camellia sinensis* (L.) O. Kuntze]

C. Lachha<sup>1\*</sup>, K. Sanjay<sup>1</sup>, M. Sandeep<sup>1</sup>, K. Vipin<sup>2</sup>, Supriya<sup>2</sup>, K. Mamta<sup>3</sup> and B. Anuradha<sup>1</sup>

<sup>1</sup>Dept. of Agronomy, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishvavidyalaya, Palampur, H.P. (176 061), India

<sup>2</sup>ICAR- National Dairy Research Institute, Karnal, Haryana (132 001), India

<sup>3</sup>Dept. of Soil Science, Rani Lakshmi Bai Central Agricultural University, Jhansi, U.P. (284 003), India

### Corresponding Author

C. Lachha

e-mail: [lachha1103@gmail.com](mailto:lachha1103@gmail.com)

### Article History

Received on 16<sup>th</sup> May, 2024

Received in revised form on 21<sup>st</sup> July, 2024

Accepted in final form on 05<sup>th</sup> August, 2024

### Abstract

A field experiment was conducted at the Research Farm of the Department of Tea Husbandry and Technology, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishvavidyalaya, Palampur (Himachal Pradesh) in pre-monsoon, monsoon and post-monsoon i.e., from March, 2021 to November, 2021. The objective was to assess the impact of organic nutrient inputs on Tea (*Camellia sinensis* L. O. Kuntze). The experiment comprised of 10 treatments with the organic treatment include application of different nitrogen sources, viz farmyard manure; vermicompost; jeevamrit and vermiwash. The experiment included ten treatments, including a control group, FYM (Farm Yard Manure) @ 20 t ha<sup>-1</sup>, vermicompost @ 10 t ha<sup>-1</sup>, split doses of FYM @ 10 t ha<sup>-1</sup>+ 10 t ha<sup>-1</sup>, split doses of vermicompost @ 5 t ha<sup>-1</sup>+5 t ha<sup>-1</sup>, FYM @ 20 t ha<sup>-1</sup>+Jeevamrit @ 10%, vermicompost @ 10 t ha<sup>-1</sup>+Jeevamrit @10%, FYM @ 20 t ha<sup>-1</sup>+vermiwash @ 10%, vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10% and Jeevamrit @ 10%. The experiment was designed following a Randomized Block Design (RBD) with three replications. The soil of experiment field was acidic in nature, silty clay loam in texture, low in Nitrogen, medium in Phosphorus and Potassium. The results showed that the growth, yield attributes and overall tea yield increased significantly with the application of vermicompost @ 10 t ha<sup>-1</sup>+Jeevamrit @ 10%. Furthermore, this treatment exhibited notably higher gross returns, net returns and additional net returns compared to the control group.

**Keywords:** FYM, Jeevamrit, vermicompost, vermiwash

### 1. Introduction

Tea, scientifically known as *Camellia sinensis* (L.) O. Kuntze, belongs to the Theaceae family and stands as one of the oldest and most significant globally consumed beverages (Jigisha et al., 2012). It is derived from the tender shoots of the tea plant, typically composed of two or three leaves and a bud (Liu et al., 2017). These shoots are harvested at intervals to produce two main types (Kumari et al., 2022): 'black' tea (which undergoes withering and fermentation) and 'green' tea (which undergoes withering but remains unfermented). Black tea, processed from the young tender shoot of fresh tea (Muthumani and Kumar, 2007). Tea is the most widely consumed beverage in the world (Hazra et al., 2019) and India holds the position of being the second-largest tea producer globally (Arya et al., 2013), following China, with an annual production of 1.4 mt, cultivated over 0.6 mha of land (Anonymous, 2020–21a). In India, tea cultivation occurs in 15 states, with Assam, West Bengal, and Kerala being the primary tea-producing regions.

The growing popularity of tea is attributed to its rich content of beneficial antioxidants, amino acids, and vitamins. As indicated in a study conducted by Expert Market Research, the Indian tea sector is projected to witness a Compound Annual Growth Rate (CAGR) of 4.2% between 2021 and 2026, with the possibility of achieving a production volume of around 1.40 mt by the conclusion of this timeframe (Ganadinni, 2021).

The tea industry in Himachal Pradesh has a history of approximately 150 years, dating back to its initiation in 1849 (Verma and Gupta, 2015). Himachal Pradesh has a total tea cultivation area of 2310.714 hectares, resulting in a tea production of 11.45 lakh kg (Anonymous, 2020–21b). In recent years, chemical farming practice has brought drastic changes in soil ecology, crop productivity and quality, in tea farming system (Shah et al., 2016; Sharma et al., 2014). Organic farming, which emphasizes the natural growth of crops without the use of chemical fertilizers, pesticides, or foreign elements, is considered a suitable remediation measure (Shahane et al., 2022; Supriya et al., 2023). Comparatively



to chemical fertilizers, organic fertilization of tea plantations may increase soil fertility and achieve carbon build-up, which was a crucial component of determining soil characteristics and production (Singh et al., 2023; Qui and Wang, 2014). The organic sources of nutrients in tea production are cheap and easily available without negative effects to the environment. Ren et al. (1999) and Sun et al. (2003) reported that application of organic fertilizers is one of important practical measures to improve soil fertility. Ji and Guo (2018) also reported that application of organic fertilizer boosts potential ecosystem function, changes the network structure, and increases soil microbial diversity. The important organic amendments are crop residues; vermicompost, farm yard manure, green manure etc. and use of these inputs are increasingly becoming important aspects of environmentally sound sustainable agriculture (Timsina and Connor, 2001). Recently, organic farming practices for tea have been gaining popularity, as compared to conventional farming, due to their effect on sustainability in tea production and improved product quality (Barbora, 1995). In Himachal Pradesh research work on systematic field experiments on supply of nutrients through organic sources, particularly on the use of vermicompost, vermiwash and *jeevamrit* in tea crop and their recommended dose is lacking. Keeping in view the above-mentioned facts the present investigation entitled “Effect of organic nutrient inputs in Tea [*Camellia sinensis* (L.) O. Kuntze]” conducted for a period of one year in 2021 at the Experimental Farm of Department of Tea husbandry and Technology, Himachal Pradesh Krishi Vishvavidyalaya, Palampur.

## 2. Materials and Methods

### 2.1. Experiment site

The field trial took place at the main campus farm of the Department of Tea Husbandry and Technology, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishvavidyalaya in Palampur, India from March-November, 2021. The experimental location is positioned at an altitude of 1291 meters above sea level, with coordinates at latitude 32°61 N and longitude 76°31 E, nestled within the Palam valley of Himachal Pradesh.

From an agricultural and climatic perspective, this experimental location represents the sub-humid temperate and mid-hill zone of Himachal Pradesh. This region is known for experiencing cool winters from November to February and mild summers from March to June. The annual precipitation varies between 1500 to 2300 mm, with approximately 70-75% of it occurring during the monsoon season from June to September. Winter rainfall typically falls during December to February, while April and May usually have minimal or no rainfall.

### 2.2. Treatments and experimental design

The experiment was carried out in a randomized block design with 10 different treatments and 3 replications. The

experiment comprised of 10 treatments viz., (1) control, (2) FYM @ 20 t ha<sup>-1</sup>, (3) vermicompost @ 10 t ha<sup>-1</sup>, (4) Split doses of FYM @ 10 t ha<sup>-1</sup>+10 t ha<sup>-1</sup>, (5) vermicompost @ 5 t ha<sup>-1</sup>+5 t ha<sup>-1</sup>, (6) FYM @ 20 t ha<sup>-1</sup>+Jeevamrit @ 10%, (7) Vermicompost @ 10 t ha<sup>-1</sup>+Jeevamrit @ 10%, (8) FYM @ 20 t ha<sup>-1</sup>+vermiwash @ 10%, (9) Vermicompost 10 t ha<sup>-1</sup>+vermiwash @ 10% and (10) Jeevamrit @ 10%.

### 2.3. Yield analysis

In each plot, a quadrat of 50×50 cm<sup>2</sup> was placed randomly at three spots. The total numbers of new growth in the form of new emerged shoots was recorded one day before each plucking. After averaging the recorded count, converted it into a value per square meter by multiplying it by a factor of 4.

Five plants from each plot were selected randomly and ten shoots per plant were tagged. Total 50 shoots plot<sup>-1</sup> were plucked for shoot biomass and the fresh weight of the 50 shoots was measured and then average weight per shoot was calculated. For dry weight per shoot, fresh shoots were sun dried for 2 days and then oven dried for 48 hrs. at 60°C and final dry weight shoot<sup>-1</sup> was calculated by taking average weight.

The two leaf and one bud from each plot were plucked manually. Made tea was prepared from the two leaves and a bud which were plucked from each plot and converted into kg ha<sup>-1</sup>

## 3. Results and Discussion

### 3.1. Number of actively growing shoots per square meter

The effect of organic nutrient sources on number of growing shoots per square meter have been presented in Table 1. The effects of organic nutrient sources on number of growing shoots per square meter were found to be significant during all three seasons. Irrespective of different treatments, number of actively growing shoots was also influenced by seasonal effect. During the monsoon season, a greater number of growing shoots per square meter were observed, followed by the pre and post monsoon seasons. Among all the treatments, a significantly higher number of growing shoots per square meter were recorded in the treatment involving the application of vermicompost @ 10 t ha<sup>-1</sup> and jeevamrit @ 10%. This result was statistically on par with all other treatments, except for the treatment involving only jeevamrit @ 10% and the control group (which did not use any organic nutrient source). A similar trend was also observed during the monsoon and post monsoon seasons. The increased number of growing shoots per square meter observed with the application of 10 t ha<sup>-1</sup> of vermicompost and jeevamrit @ 10% could be attributed to the relatively higher quantity of organic matter added. This organic matter likely improved the biological properties of the soil, leading to increased nutrient availability for the crop through mineralization and solubilization processes. Organic matter enriches soil by serving as a nutrient storehouse, guarding against nutrient



Table 1: Effect of different organic nutrient sources on number of actively growing shoots per square meter

Treatments details		No. of actively growing shoots m <sup>-2</sup>		
		Pre-monsoon season	Monsoon season	Post monsoon season
T <sub>1</sub>	Control	183	199	174
T <sub>2</sub>	FYM @ 20 t ha <sup>-1</sup>	201	216	190
T <sub>3</sub>	Vermicompost @ 10 t ha <sup>-1</sup>	208	222	198
T <sub>4</sub>	FYM @ 10 t ha <sup>-1</sup> (pre-monsoon season)+@ 10 t ha <sup>-1</sup> (monsoon season)	200	218	193
T <sub>5</sub>	Vermicompost @ 5 t ha <sup>-1</sup> (pre-monsoon season) +@ 5 t ha <sup>-1</sup> (monsoon season)	201	223	200
T <sub>6</sub>	FYM @ 20 t ha <sup>-1</sup> + Jeevamrit @ 10% within 2 day after plucking	207	220	194
T <sub>7</sub>	Vermicompost @ 10 t ha <sup>-1</sup> +Jeevamrit @ 10% within 2 day after plucking	212	228	203
T <sub>8</sub>	FYM @ 20 t ha <sup>-1</sup> + Vermiwash @ 10% within 2 days after plucking	203	219	194
T <sub>9</sub>	Vermicompost @ 10 t ha <sup>-1</sup> + Vermiwash @ 10% within 2 days after plucking	210	226	202
T <sub>10</sub>	Jeevamrit @ 10% within 2 days after plucking	186	201	177
SEm±		5.67	5.91	5.38
CD (p=0.05)		16.64	17.35	15.77

loss, and aiding in nutrient delivery to plants. It stimulates microbial activity, breaking down organic substances into forms accessible to plants. Moreover, its elevated cation exchange capacity (CEC) enables it to draw in and gradually dispense crucial nutrients, guaranteeing a consistent nutrient source for plant development. These results are in close conformity with Negi et al. (2017), Islama et al. (2017).

### 3.2. Mean dry shoot biomass (mg shoot<sup>-1</sup>)

The data embodied in Table 2 revealed that the effect of organic nutrient sources on dry shoot biomass (mg shoot<sup>-1</sup>) was found to be significant in all the three seasons. Among all the treatments, the treatment comprising vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% recorded maximum dry shoot biomass in premonsoon season which was statistically at par with vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%. Similar trend

Table 2: Effect of different organic nutrient sources on mean dry shoot biomass

T. N.	Treatments details	Dry shoot biomass (mg shoot <sup>-1</sup> )		
		Pre-monsoon season	Monsoon season	Post monsoon season
T <sub>1</sub>	Control	44.0	45.0	40.0
T <sub>2</sub>	FYM @ 20 t ha <sup>-1</sup>	54.0	53.0	49.0
T <sub>3</sub>	Vermicompost @ 10 t ha <sup>-1</sup>	55.0	54.0	51.0
T <sub>4</sub>	FYM @ 10 t ha <sup>-1</sup> (pre-monsoon season)+@ 10 t ha <sup>-1</sup> (monsoon season)	49.0	49.7	45.0
T <sub>5</sub>	Vermicompost @ 5 t ha <sup>-1</sup> (pre-monsoon season) +@ 5 t ha <sup>-1</sup> (monsoon season)	51.0	51.0	47.0
T <sub>6</sub>	FYM @ 20 t ha <sup>-1</sup> + Jeevamrit @ 10% within 2 day after plucking	58.0	57.0	54.0
T <sub>7</sub>	Vermicompost @ 10 t ha <sup>-1</sup> +Jeevamrit @ 10% within 2 day after plucking	63.0	60.0	58.0
T <sub>8</sub>	FYM @ 20 t ha <sup>-1</sup> + Vermiwash @ 10% within 2 days after plucking	58.0	56.0	53.0
T <sub>9</sub>	Vermicompost @ 10 t ha <sup>-1</sup> + Vermiwash @ 10% within 2 days after plucking	60.0	58.0	55.0
T <sub>10</sub>	Jeevamrit @ 10% within 2 days after plucking	46.0	47.0	42.0
SEm±		1.6	1.6	1.2
CD (p=0.05)		4.8	4.6	3.6



was also followed in post monsoon season. Significantly higher dry biomass was recorded with application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% in monsoon season which was statistically at par with treatments having vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%, FYM @ 20 t ha<sup>-1</sup>+jeevamrit @ 10% and FYM @ 20 t ha<sup>-1</sup>+vermiwash @ 10%. Dry shoot biomass was higher in treatment vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% due to increased microbial activity which might have resulted into steady supply of nutrients, which ultimately increased the dry shoot biomass. Organic matter enhances nutrient availability in soil by acting as a reservoir, preventing nutrient leaching, and facilitating nutrient release to plants. It fosters microbial activity, breaking down organic materials into plant-accessible forms. Additionally, its high cation exchange capacity (CEC) allows it to attract and slowly release essential nutrients, ensuring a steady supply for plant growth. These results are in close conformity with Negi and Bisht (2017).

### 3.3. Made black tea

The effects of organic nutrient sources on made black tea yield (Figure 1) was found to be significant during all three seasons. In pre-monsoon season among all the treatments, higher made black tea yield was observed with the application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @10% which was statistically at par with the application vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%. The lowest made black tea yield was recorded in absolute control. Similar trend was also followed in monsoon and post monsoon season. These results are in close conformity with Negi and Bisht (2017).




Figure 1: Effect of different organic nutrient sources on made black tea yield

### 3.4. Economics

The economics of crop production in terms of net return, additional net return over control and net return per rupee invested have greater impact on practical utility and acceptance of any technology. In order to evaluate the economic feasibility of different organic treatments, cost of cultivation, gross return, and additional net return over control and net return per rupee invested were worked out and the results have been depicted graphically in Table 3.

Table 3: Effect of different organic nutrient sources on economics of made black tea

T. N.	Gross return	Net return	Additional net return over control	Net return per rupee invested
T <sub>1</sub>	354053.3	202235.6	-	1.33
T <sub>2</sub>	474160.4	267506.9	65271.33	1.29
T <sub>3</sub>	528880	300297.8	98062.18	1.31
T <sub>4</sub>	465661.8	262152.3	59916.7	1.29
T <sub>5</sub>	523893.3	297217.8	94982.18	1.31
T <sub>6</sub>	500964.7	283835	81599.39	1.31
T <sub>7</sub>	562320	321524.4	119288.8	1.34
T <sub>8</sub>	492009.8	282613.9	80378.33	1.35
T <sub>9</sub>	542813.3	309024.4	106788.8	1.32
T <sub>10</sub>	380160	218135.6	15899.96	1.35

#### 3.4.1. Cost of cultivation

The highest cost of cultivation (240795.6 ₹ ha<sup>-1</sup>) was incurred with application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% as compared to all the remaining treatments. The least cost was incurred in control. The higher cost in vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% could be attributed to the use of multiple inputs like vermicompost and jeevamrit in the treatment.

#### 3.4.2. Gross returns

The application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% recorded highest gross return of ₹ 562320 ha<sup>-1</sup> which was followed by vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%. The lowest gross return was recorded in absolute control. The significant difference in gross return was mainly because of the difference in made tea yield due to treatment effect.

#### 3.4.3. Net return

The application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% recorded significantly higher net return of ₹ 562320 ha<sup>-1</sup> which was followed by vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%. The lowest net return was recorded in absolute control. The significant difference in net return was mainly because of the difference in made tea yield and cost of cultivation.

#### 3.4.4. Additional net return over control

The application of vermicompost @ 10 t ha<sup>-1</sup>+jeevamrit @ 10% recorded higher additional net return of ₹ 119288 ha<sup>-1</sup> which was followed by vermicompost @ 10 t ha<sup>-1</sup>+vermiwash @ 10%.

#### 3.4.5. Net return per rupee invested

The net profit for each rupee invested was notably influenced by various organic treatments. Specifically, the application of 20 t ha<sup>-1</sup> of Farm Yard Manure (FYM) in combination with 10% jeevamrit yielded the highest net return per rupee invested. This superior ratio in net return per rupee invested in the



treatment using FYM at 20 t ha<sup>-1</sup> and 10% jeevamrit was attributed to the comparatively lower input costs."

#### 4. Conclusion

The Application of organic fertilizer enhanced plant height, number of leaves and mean fresh and dry biomass. Application of vermicompost @ 10 t ha<sup>-1</sup> followed by foliar application of jeevamrit @ 10% gave higher green leaf yield as well as black made tea yield while the highest values of economic indices was recorded in treatment having vermicompost @ 10 t ha<sup>-1</sup> +Jeevamrit @ 10%.

#### 6. References

Anonymous, 2020a. Economic Survey 2020–21. Statistical Appendix, Government of India 2, 34-35. <https://www.indiabudget.gov.in/economic-survey/statistical-app-volume-2>.

Anonymous, 2020b. Economic Survey 2021–22. Economic and Statistical Department, Government of Himachal Pradesh 2, 89.

Arya, N., 2013. Indian tea scenario. International Journal of Scientific and Research Publications 3(7), 1–10.

Barbora, A.C., 1995. Sulphur management for tea in north eastern India. Sulphur in Agriculture 19, 9–15.

Ganadinni, S., 2021. Trends and future of tea trade in India and the role of supply chain. <https://www.indianretailer.com/article/whats-hot/retail-trends/trends-and-future-of-tea-trade-in-india-and-the-role-of-supply-chain>.

Hazra, A., Dasgupta, N., Sengupta, C., Bera, B., Das, S., 2019. Tea: A worthwhile, popular beverage crop since time immemorial. Agronomic Crops: Volume 1: Production Technologies, 507–531.

Islama, S., Hamid, F.S., Aminc, K., Sumreend, S., Zamane, Q., Khanf, N., Khang, A., Shahh, B.S., 2017. Effect of organic fertilizer on the growth of Tea (*Camellia sinensis* L.). International Journal of Sciences: Basic and Applied Research 36(8), 1–9.

Ji, L., Wu, Z., You, Z., Yi, X., Ni, K., Guo, S., Ruan, J., 2018. Effects of organic substitution for synthetic N fertilizer on soil bacterial diversity and community composition: a 10-year field trial in a tea plantation. Agriculture, Ecosystems and Environment 268, 124–132.

Jigisha, A., Nishant, R., Navin, K., Pankaj, G., 2012. Green tea: A magical herb with miraculous outcomes. International Research Journal of Pharmacy 3(5), 139–148.

Kumari, A., Kumar, D., 2022. Evaluation of antioxidant and cytotoxic activity of herbal teas from Western Himalayan region: a comparison with green tea (*Camellia sinensis*) and black tea. Chemical and Biological Technologies in Agriculture 9(1), 33.

Liu, S., Gao, J., Chen, Z., Qiao, X., Huang, H., Cui, B., Liu, J., 2017. Comparative proteomics reveals the physiological

differences between winter tender shoots and spring tender shoots of a novel tea (*Camellia sinensis* L.) cultivar ever-growing in winter. BMC Plant Biology 17, 1–12.

Muthumani, T., Kumar, R.S., 2007. Studies on freeze-withering in black tea manufacturing. Food Chemistry 101(1), 103–106.

Negi, G.C.S., Bisht, V., 2017. Promoting organic tea farming in mid-hills of north-west Himalaya, India. Tea 38(2), 57–67.

Qui, S.L., Wang, L.M., Huang, Lin, X.J., 2014. Effects of fertilization regimes on tea yields, soil fertility, and soil microbial diversity. Chilean Journal of Agricultural Research 74 (3), 333–339.

Ren, Z.G., Chen, Y.S., Tang, F.Q., 1999. Effect of inorganic fertilizer combined with organic manure on the micro flora and enzyme activities in paddy soil. Plant Nutrition and Fertilizer Science 2, 279–283.

Shah, S.K., Pate, V.A., 2016. Tea production in India: challenges and opportunities. Journal of Tea Science Research 6, 23–25.

Shahane, A.A., Shivay, Y.S., 2022. Soil health management in organic production system-A review. International Journal of Bio-resource and Stress Management 13(11), 1186–1200.

Sharma, A., Sharma, R.P., Sharma, G.D., Sankhyan, N.K., Sharma, M., 2014. Integrated nutrient supply system for cauliflower French bean- Okra cropping sequence in humid temperate zone of North Western Himalayas. Indian Journal of Horticulture 71, 211–216.

Singh, A.K., Chauhan, R.K., Kumar, C., Bera, B., 2023. Sustaining Darjeeling organic tea (*Camellia sinensis* L) production and tea quality through organic inputs in the Himalayan foothills. Journal of Soil Science and Plant Nutrition 23(2), 1774–1796.

Sun, R.L., Zhao, B.Q., Zhu, L.S., 2003. Effects of long-term fertilization on soil enzyme activities and its role in adjusting controlling soil fertility. Plant Nutrition and Fertilizer Science 9, 406–410.

Supriya, Singh, D.K., Ram, N., Kumar, R.K.M., Sharma, U., Kumar, B., Verma, P., 2023. Comparative analysis of traditional scented rice (*Oryza sativa* L.) varieties under organic cultivation based on DUS testing and yield assessment. International Journal of Plant & Soil Science 35(18), 1474–1483.

Timsina, J., Connor, D.J., 2001. Productivity and management of rice-wheat systems: Issues and challenges. Field Crops Research 69, 93–132.

Verma, P., Gupta, S., 2015. Factors affecting socio-economic status of farm workers of tea industry in Himachal Pradesh. Himachal Journal of Agricultural Research 41(1), 36–41.

